



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 239/82, 487/22, 487/14, 487/04, A61K 31/517, 31/519, 31/496	A1	(11) International Publication Number: WO 00/43374 (43) International Publication Date: 27 July 2000 (27.07.00)
(21) International Application Number: PCT/US00/01775 (22) International Filing Date: 24 January 2000 (24.01.00) (30) Priority Date: 60/117,255 26 January 1999 (26.01.99) US (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BOCK, Mark, G. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). PATANE, Michael, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). STEELE, Thomas, G. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GI, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: POLYAZANAPHTHALENONE DERIVATIVES USEFUL AS ALPHA 1A ADRENOCEPTOR ANTAGONISTS (57) Abstract <p>Dihydro-polyazanaphthalen-2-one compounds (e.g., dihydroquinazolin-2-one and dihydropteridin-2-one derivatives) and pharmaceutically acceptable salts thereof are disclosed. The synthesis of these compounds and their use as alpha 1a adrenergic receptor antagonists is also described. One application of these compounds is in the treatment of benign prostatic hyperplasia. These compounds are selective in their ability to relax smooth muscle tissue enriched in the alpha 1a receptor subtype without at the same time inducing hypotension. One such tissue is found surrounding the urethral lining. Therefore, one utility of the instant compounds is to provide acute relief to males suffering from benign prostatic hyperplasia, by permitting less hindered urine flow. Another utility of the instant compounds is provided by combination with a human 5-alpha reductase inhibitory compound, such that both acute and chronic relief from the effects of benign prostatic hyperplasia can be achieved.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TITLE OF THE INVENTION
POLYAZANAPHTHALENONE DERIVATIVES USEFUL AS ALPHA 1A
ADRENOCEPTOR ANTAGONISTS

5 FIELD OF THE INVENTION

This invention relates to polyazananaphthalen-2-one derivatives (e.g., quinazolin-2-one and pteridin-2-one derivatives) and pharmaceutically acceptable salts thereof, their synthesis, and their use as alpha 1a adrenoceptor antagonists. More particularly, the compounds of the present invention are useful for treating benign
10 prostatic hyperplasia (BPH).

References are made throughout this application to various publications, the disclosures of which are hereby incorporated by reference in their entireties, in order to more fully describe the state of the art to which this invention
15 pertains.

BACKGROUND OF THE INVENTION

Human adrenergic receptors are integral membrane proteins which have been classified into two broad classes, the alpha and the beta adrenergic receptors. Both types mediate the action of the peripheral sympathetic nervous system
20 upon binding of catecholamines, norepinephrine and epinephrine.

Norepinephrine is produced by adrenergic nerve endings, while epinephrine is produced by the adrenal medulla. The binding affinity of adrenergic receptors for these compounds forms one basis of the classification: alpha receptors bind norepinephrine more strongly than epinephrine and much more strongly than the
25 synthetic compound isoproterenol. The binding affinity of these hormones is reversed for the beta receptors. In many tissues, the functional responses, such as smooth muscle contraction, induced by alpha receptor activation are opposed to responses induced by beta receptor binding.

Subsequently, the functional distinction between alpha and beta
30 receptors was further highlighted and refined by the pharmacological characterization of these receptors from various animal and tissue sources. As a result, alpha and beta adrenergic receptors were further subdivided into alpha 1, alpha 2, β_1 , and β_2 subtypes. Functional differences between alpha 1 and alpha 2 receptors have been recognized, and compounds which exhibit selective binding between these two
35 subtypes have been developed.

For a general background on the alpha adrenergic receptors, the reader's attention is directed to Robert R. Ruffolo, Jr., α -Adrenoreceptors: Molecular Biology, Biochemistry and Pharmacology, (Progress in Basic and Clinical Pharmacology series, Karger, 1991), wherein the basis of alpha 1/alpha 2 subclassification, the molecular biology, signal transduction (G-protein interaction and location of the significant site for this and ligand binding activity away from the 3'-terminus of alpha adrenergic receptors), agonist structure-activity relationships, receptor functions, and therapeutic applications for compounds exhibiting alpha-adrenergic receptor affinity was explored.

The cloning, sequencing and expression of alpha receptor subtypes from animal tissues has led to the subclassification of the alpha 1 receptors into alpha 1d (formerly known as alpha 1a or 1a/1d), alpha 1b and alpha 1a (formerly known as alpha 1c) subtypes. Each alpha 1 receptor subtype exhibits its own pharmacologic and tissue specificities. The designation "alpha 1a" is the appellation recently approved by the IUPHAR Nomenclature Committee for the previously designated "alpha 1c" cloned subtype as outlined in the 1995 Receptor and Ion Channel Nomenclature Supplement (Watson and Girdlestone, 1995). The designation alpha 1a is used throughout this application to refer to this subtype. At the same time, the receptor formerly designated alpha 1a was renamed alpha 1d. The new nomenclature is used throughout this application. Stable cell lines expressing these alpha 1 receptor subtypes are referred to herein; however, these cell lines were deposited with the American Type Culture Collection (ATCC) under the old nomenclature. For a review of the classification of alpha 1 adrenoceptor subtypes, see, Michel et al., *Naunyn-Schmiedeberg's Arch. Pharmacol.* (1995), 352:1-10.

The differences in the alpha adrenergic receptor subtypes have relevance in pathophysiological conditions. Benign prostatic hyperplasia, also known as benign prostatic hypertrophy or BPH, is an illness typically affecting men over fifty years of age, increasing in severity with increasing age. The symptoms of the condition include, but are not limited to, increased difficulty in urination and sexual dysfunction. These symptoms are induced by enlargement, or hyperplasia, of the prostate gland. As the prostate increases in size, it impinges on free-flow of fluids through the male urethra. Concomitantly, the increased noradrenergic innervation of the enlarged prostate leads to an increased adrenergic tone of the bladder neck and urethra, further restricting the flow of urine through the urethra.

In benign prostatic hyperplasia, the male hormone 5 α -dihydrotestosterone has been identified as the principal culprit. The continual production of 5 α -dihydrotestosterone by the male testes induces incremental growth of the prostate gland throughout the life of the male. Beyond the age of about fifty years, in many men, this enlarged gland begins to obstruct the urethra with the pathologic symptoms noted above.

The elucidation of the mechanism summarized above has resulted in the recent development of effective agents to control, and in many cases reverse, the pernicious advance of BPH. In the forefront of these agents is Merck & Co., Inc.'s product PROSCAR® (finasteride). The effect of this compound is to inhibit the enzyme testosterone 5- α reductase, which converts testosterone into 5 α -dihydrotestosterone, resulting in a reduced rate of prostatic enlargement, and often reduction in prostatic mass.

The development of such agents as PROSCAR® bodes well for the long-term control of BPH. However, as may be appreciated from the lengthy development of the syndrome, its reversal also is not immediate. In the interim, those males suffering with BPH continue to suffer, and may in fact lose hope that the agents are working sufficiently rapidly.

In response to this problem, one solution is to identify pharmaceutically active compounds which complement slower-acting therapeutics by providing acute relief. Agents which induce relaxation of the lower urinary tract tissue, by binding to alpha 1 adrenergic receptors, thus reducing the increased adrenergic tone due to the disease, would be good candidates for this activity. Thus, one such agent is alfuzosin, which is reported in EP 0 204597 to induce urination in cases of prostatic hyperplasia. Likewise, in WO 92/00073, the selective ability of the R(+) enantiomer of terazosin to bind to adrenergic receptors of the alpha 1 subtype was reported. In addition, in WO 92/16213, combinations of 5 α -reductase inhibitory compounds and alpha1-adrenergic receptor blockers (terazosin, doxazosin, prazosin, bunazosin, indoramin, alfuzosin) were disclosed. However, no information as to the alpha 1d, alpha 1b, or alpha 1a subtype specificity of these compounds was provided as this data and its relevancy to the treatment of BPH was not known. Current therapy for BPH uses existing non-selective alpha 1 antagonists such as prazosin (Minipress, Pfizer), Terazosin (Hytrin, Abbott) or doxazosin mesylate (Cardura, Pfizer). These non-selective antagonists suffer from side effects related to antagonism of the alpha 1d and alpha 1b receptors in the peripheral vasculature, e.g., hypotension and syncope.

The relatively recent cloning of the human alpha 1a adrenergic receptor (ATCC CRL 11140) and the use of a screening assay utilizing the cloned human alpha 1a receptor has enabled identification of compounds which specifically interact with the human alpha 1a adrenergic receptor. For further description, see WO 94/08040 and WO 94/10989. As disclosed in the instant patent disclosure, a cloned human
5 alpha 1a adrenergic receptor and a method for identifying compounds which bind the human alpha 1a receptor has made possible the identification of selective human alpha 1a adrenergic receptor antagonists useful for treating BPH.

Several classes of compounds have been disclosed to be selective alpha
10 1a adrenergic receptor antagonists useful for treating BPH. WO 94/22829 discloses, for example, certain 4-(un)substituted phenyl-1,4-dihydropyridine derivatives which are described as potent, selective alpha 1a antagonists with weak calcium channel antagonistic activity and which are further described to be anticipated as useful for treating BPH. As another example, WO 96/14846, WO 97/17969 and WO 97/42956
15 each disclose certain dihydropyrimidine derivatives (e.g., certain 1,2,3,6-tetrahydro-2-oxo-pyrimidine derivatives) which are selective antagonists for the human alpha 1a receptor and useful for treatment of BPH, impotency, cardiac arrhythmia, and other diseases where antagonism of the alpha 1a receptor may be useful. As still another example, WO 96/40135 discloses, *inter alia*, certain phenylpiperidinyl alkyl saccharin
20 derivatives and their use as selective alpha 1a antagonists. Other examples are US 5661163 and WO 96/40136, which disclose, *inter alia*, piperidinyl- and piperazinyl-alkyl-substituted phenyl acetamides. Yet another example is EP 748840, which discloses, *inter alia*, certain arylpiperazinylpropyl substituted pyrimidinediones useful as alpha 1 adrenoceptor antagonists. Still other alpha 1a selective antagonist
25 compounds are disclosed in WO 98/57632, WO 98/57638, WO 98/57639, WO 98/57640, WO 98/57641 and WO 98/57642.

The instant patent disclosure discloses novel dihydro-polyazaphthalen-2-one compounds which selectively bind to the human alpha 1a
30 receptor. These compounds are further tested for binding to other human alpha 1 receptor subtypes, as well as counterscreened against other types of receptors (e.g., alpha 2), thus defining the specificity of the compounds of the present invention for the human alpha 1a adrenergic receptor.

It is an object of the present invention to identify compounds which bind to the alpha 1a adrenergic receptor. It is a further object of the invention to
35 identify compounds which act as antagonists of the alpha 1a adrenergic receptor. It is


another object of the invention to identify alpha 1a adrenergic receptor antagonist compounds which are useful agents for treating BPH in animals, preferably mammals, especially humans. Still another object of the invention is to identify alpha 1a adrenergic receptor antagonists which are useful for relaxing lower urinary tract tissue in animals, preferably mammals, especially humans.

The compounds of the present invention are alpha 1a adrenergic receptor antagonists. Thus, the compounds of the present invention are useful for treating BPH in mammals. Additionally, it has been found that the alpha 1a adrenergic receptor antagonists of the present invention are also useful for relaxing lower urinary tract tissue in mammals.

SUMMARY OF THE INVENTION

The present invention provides dihydro-polyazananaphthalen-2-one compounds and pharmaceutically acceptable salts thereof for the treatment of urinary obstruction caused by benign prostatic hyperplasia (BPH). The compounds antagonize the human alpha 1a adrenergic receptor at nanomolar and subnanomolar concentrations while exhibiting lower affinity for the alpha 1d and alpha 1b human adrenergic receptors and many other G-protein coupled receptors. This invention has the advantage over non-selective alpha 1 adrenoceptor antagonists of reduced side effects related to peripheral adrenergic blockade. Such side effects include hypotension, syncope, lethargy, etc.

More particularly, the present invention is a compound of formula (I):



wherein Q is selected from the group consisting of:

A¹, A², A³ and A⁴ are each independently selected from C-X² and N, provided that no more than two of A¹, A², A³ and A⁴ are N, and further provided that when two of A¹, A², A³ and A⁴ are N, each nitrogen atom is bonded to two carbon atoms;

- 5 each X¹ is independently hydrogen, halogen, cyano, nitro, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl, or (CH₂)₀₋₄OR^a;

each X² is independently hydrogen, halogen, cyano, nitro, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl, or (CH₂)₀₋₄OR^a;

10

R¹ is hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl, phenyl, or substituted phenyl, wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, nitro, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl and (CH₂)₀₋₄OR^a;

15

R² is hydrogen, C₁-C₈ alkyl, or fluorinated C₁-C₈ alkyl;

R³ is hydrogen, C₁-C₈ alkyl, or fluorinated C₁-C₈ alkyl;

20

Y is carbon or nitrogen, provided that when Y is nitrogen, R⁵ is absent;

Z is CH₂, CHOR^b, CHF, CHR^b, C(R^b)₂, CF₂, CHCHF₂, C=CF₂, or C=O;

- 25 E, G, L and M are each independently selected from hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, (CH₂)₀₋₄OR^c, (CH₂)₀₋₄N(R^c)₂, (CH₂)₀₋₄CN, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂R^c and (CH₂)₀₋₄SO₂N(R^c)₂;

- 30 J is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, (CH₂)₁₋₄OR^c, (CH₂)₁₋₄N(R^c)₂, (CH₂)₁₋₄CN, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂R^c, or (CH₂)₀₋₄SO₂N(R^c)₂;

- R^4 is phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, thienyl, furanyl, substituted pyridyl, substituted thienyl, or substituted furanyl; wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, OR^c , $(CH_2)_0-3CON(R^c)_2$, $(CH_2)_0-3CO_2R^c$, methylenedioxy when the phenyl ring is di-substituted and the substituents are on adjacent carbon atoms, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; and wherein the substituents on the substituted naphthyl, pyridyl, thienyl, or furanyl are independently selected from phenyl, OR^c , halogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl, C₃-C₈ cycloalkyl and fluorinated C₃-C₈ cycloalkyl;
- R^5 is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, tetrazole, isooxadiazole, phenyl, or substituted phenyl; wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, $(CH_2)_0-3CON(R^c)_2$, $(CH_2)_0-3CO_2R^c$, methylenedioxy when the phenyl ring is di-substituted and the substituents are on adjacent carbon atoms, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;
- R^6 and R^7 are each independently selected from hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl and fluorinated C₃-C₈ cycloalkyl;
- R^8 is phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, substituted pyridyl, pyridyl N-oxide ($N\textcircled{O}$), substituted pyridyl N-oxide, pyrazinyl, substituted pyrazinyl, thienyl, substituted thienyl, thiazolyl, substituted thiazolyl, furanyl, substituted furanyl, quinazolinyl, or substituted quinazolinyl; wherein the substituents on the phenyl are independently selected from halogen, cyano, nitro, OR^c , $N(R^c)_2$, NR^cCOR^c , $NR^cCON(R^c)_2$, $NR^cSO_2R^c$, $NR^cSO_2N(R^c)_2$, $(CH_2)_0-4CO_2R^c$, $(CH_2)_0-4CON(R^c)_2$, $(CH_2)_0-4SO_2N(R^c)_2$, $(CH_2)_0-4SO_2R^c$, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; and wherein the substituents on the substituted naphthyl, pyridyl, pyrazinyl, thienyl, thiazolyl, furanyl, or quinazolinyl are independently selected from cyano, nitro, $N(R^c)_2$, $(CH_2)_0-4CO_2R^c$, $(CH_2)_0-4CON(R^c)_2$, $(CH_2)_0-4SO_2N(R^c)_2$, $(CH_2)_0-4SO_2R^c$, phenyl, OR^c , halogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl, C₃-C₈ cycloalkyl and fluorinated C₃-C₈ cycloalkyl;
- R^9 is hydrogen, C₁-C₈ alkyl, or fluorinated C₁-C₈ alkyl;

- R^{10} is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, tetrazole, isooxadiazole, phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, substituted pyridyl, thienyl, substituted thienyl, furanyl, or substituted furanyl; wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, nitro, OR^c , $(CH_2)_0-4CO_2R^c$, $(CH_2)_0-4CON(R^c)_2$, $N(R^c)_2$, NR^cCOR^c , $NR^cCON(R^c)_2$, $NR^cSO_2R^c$, $NR^cSO_2N(R^c)_2$, $(CH_2)_0-4SO_2N(R^c)_2$, $(CH_2)_0-4SO_2R^c$, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; and wherein the substituents on the substituted naphthyl, pyridyl, thienyl, or furanyl are independently selected from $(CH_2)_0-4CO_2R^c$, $(CH_2)_0-4CON(R^c)_2$, $(CH_2)_0-4SO_2N(R^c)_2$, $(CH_2)_0-4SO_2R^c$, phenyl, OR^c , halogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl and C₃-C₈ cycloalkyl;
- R^{13} , R^{14} , R^{15} and R^{16} are each independently selected from hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, and $(CH_2)_2-4OR^c$;
- R^{18} and R^{20} are each independently selected from hydrogen and OR^d ;
- R^a is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, or C₃-C₈ cycloalkyl;
- R^b is C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or fluorinated C₃-C₈ cycloalkyl;
- R^c is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or fluorinated C₃-C₈ cycloalkyl;
- R^d is hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or $(CH_2)_0-4CF_3$;
- m, n, o, and p are each independently integers from 0 to 3;
- q₁ is an integer from 0 to 5;
- t is an integer from 2 to 5;

u and v are each independently integers from 0 to 3; provided that u and v are not both zero; and further provided that when u = 0, Z is selected from CH₂, CHF, CHR^b, C(R^b)₂, CF₂, CHCHF₂ and C=CF₂; and

5 w is an integer from 0 to 3, provided that when w is 0, R²⁰ is hydrogen;

or a pharmaceutically acceptable salt thereof.

The present invention also includes pharmaceutical compositions, methods of preparing pharmaceutical compositions, and methods of treatment.

10 These and other embodiments, aspects and features of the present invention are either further described in or will be apparent from the ensuing description, examples and appended claims.

DETAILED DESCRIPTION OF THE INVENTION

15 The present invention includes dihydro-polyazanaphthalen-2-one compounds of Formula (I) above. These compounds and their pharmaceutically acceptable salts are useful as alpha 1a antagonists.

In a first embodiment, the present invention is a compound of Formula (I), wherein

20 each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, or (CH₂)₀₋₄OR^a;

25 each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, or (CH₂)₀₋₄OR^a;

R¹ is hydrogen, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, phenyl, or substituted phenyl, wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, C₁-C₄ alkyl, 30 fluorinated C₁-C₄ alkyl and (CH₂)₀₋₄OR^a;

R² is hydrogen or C₁-C₄ alkyl;

R³ is hydrogen or C₁-C₄ alkyl;

E, G, L and M are each independently selected from hydrogen and C₁-C₄ alkyl;

J is hydrogen or C₁-C₄ alkyl;

5

R⁴ is phenyl, mono- or di- or tri-substituted phenyl, pyridyl, thienyl, or furanyl; wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, CO₂R^c, OR^c, CON(R^c)₂, methylenedioxy, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

10

R⁵ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, CO₂R^c, OR^c, CON(R^c)₂, methylenedioxy, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

15

one of R⁶ and R⁷ is hydrogen, and the other of R⁶ and R⁷ is hydrogen or C₁-C₄ alkyl;

20 R⁸ is phenyl, mono- or di- or tri-substituted phenyl, pyridyl, or mono- or di- or tri-substituted pyridyl; wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, C₁₋₄ alkyl and CF₃; and wherein the substituents on the substituted pyridyl are independently selected from cyano, nitro, N(R^c)₂, (CH₂)₀₋₄CO₂R^c,
25 (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, phenyl, OR^c, halogen, C₁-C₄ alkyl and CF₃;

R⁹ is hydrogen, C₁-C₄ alkyl, or fluorinated C₁-C₄ alkyl;

30

R¹⁰ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c and C₁-C₄ alkyl;

R¹³, R¹⁴, R¹⁵ and R¹⁶ are each independently selected from hydrogen and C₁-C₄ alkyl;

5 R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^b is C₁-C₄ alkyl or (CH₂)₀₋₃CF₃;

R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

10 R^d is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

m, n, o, and p are each independently 0 or 1;

15 q_l is an integer from 0 to 4; and

all other variables are as originally defined above;

or a pharmaceutically acceptable salt thereof.

20

In a second embodiment, the present invention is a compound of Formula (I),

wherein each of A¹, A², A³ and A⁴ is C-X²; and


25

all other variables are as defined in the first embodiment

or a pharmaceutically acceptable salt thereof.

30

In a third embodiment, the present invention is a compound of Formula (I), wherein Q is



each of A¹, A², A³ and A⁴ is C-X²;

- 5 each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, or (CH₂)₀₋₄OR^a;

each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, or (CH₂)₀₋₄OR^a;

10

R¹ is hydrogen, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, phenyl, or substituted phenyl, wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl and (CH₂)₀₋₄OR^a;

15

R² is hydrogen or C₁-C₄ alkyl;

R³ is hydrogen or C₁-C₄ alkyl;

- 20 R⁴ is phenyl, mono- or di- or tri-substituted phenyl, pyridyl, thienyl, or furanyl; wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, CO₂R^c, OR^c, CON(R^c)₂, methylenedioxy, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

- 25 R⁵ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, CO₂R^c, OR^c, CON(R^c)₂, methylenedioxy, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

one of R^6 and R^7 is hydrogen, and the other of R^6 and R^7 is hydrogen or C₁-C₄ alkyl;

R^a is hydrogen, C₁-C₄ alkyl, or $(CH_2)_0-3CF_3$;

5

R^b is C₁-C₄ alkyl or $(CH_2)_0-3CF_3$;

R^c is hydrogen, C₁-C₄ alkyl, or $(CH_2)_0-3CF_3$;


10 q_1 is an integer from 0 to 4; and

all other variables are as originally defined above;

or a pharmaceutically acceptable salt thereof.

15

In a first class of the invention is a compound of Formula (II):



wherein Z is CH₂, CHOH or C=O;

20

each X^3 is independently hydrogen, CF₃, cyano, halogen, or C₁-C₄ alkyl;

R^5 is hydrogen, cyano, or CO₂R^c;


25 q_2 is an integer from 0 to 4;

q_3 is an integer from 0 to 3; and

all other variables are as defined in the third embodiment;

5 or a pharmaceutically acceptable salt thereof.

In a sub-class of the first class is a compound of formula (III):



10

each X^1 is independently hydrogen, fluorine, methyl, ethyl, cyano, CF_3 , methoxy, ethoxy, or OCF_3 ;

each X^3 is independently hydrogen, CF_3 , cyano, fluorine, methyl, or ethyl;

15

q_1 is an integer from 0 to 2; and

all other variables are as defined in the first class;

20 or a pharmaceutically acceptable salt thereof.

Exemplifying the invention is a compound selected from the group consisting of:

E, G, L and M are each independently selected from hydrogen and C₁-C₄ alkyl;

J is hydrogen or C₁-C₄ alkyl;

5

one of R⁶ and R⁷ is hydrogen, and the other of R⁶ and R⁷ is hydrogen or C₁-C₄ alkyl;

10 R⁸ is selected from phenyl, mono- or di- or tri-substituted phenyl, pyridyl, or mono- or di- or tri-substituted pyridyl; wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, C₁₋₄ alkyl and CF₃; and wherein the substituents on the substituted pyridyl are independently selected from
15 cyano, N(R^c)₂, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, phenyl, OR^c, halogen, C₁-C₄ alkyl and CF₃;

20 R¹⁰ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c and C₁-C₄ alkyl;

25 R¹³, R¹⁴, R¹⁵ and R¹⁶ are each independently selected from hydrogen and C₁-C₄ alkyl;

R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

30

R^d is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₄CF₃;


m, n, o, and p are each independently 0 or 1;

q1 is an integer from 0 to 4; and

all other variables are as originally defined above;

- 5 or a pharmaceutically acceptable salt thereof.

In a second class of the invention is a compound of Formula (IV):



- 10 wherein T is C-X³ or N;

each X³ is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, C₁₋₄ alkyl, or CF₃;

- 15 R¹⁰ is hydrogen, cyano, or OR^c;

R²⁰ is hydrogen or OH;

q2 is an integer from 0 to 4;

20

q3 is an integer from 0 to 3;

w is 0 or 1, provided that when w is 0, R²⁰ is hydrogen; and

- 25 all other variables are as defined in the fourth embodiment;

or a pharmaceutically acceptable salt thereof.

In a sub-class of the second class is a compound of formula (IV),
wherein each X¹ is fluorine;

5 each X² is hydrogen;

R²⁰ is hydrogen;

q₁ is an integer from 0 to 3;

10

w is 1; and

all other variables are as defined in the second class;

15 or a pharmaceutically acceptable salt thereof.

Also exemplifying the invention is a compound selected from the
group consisting of:

20 *trans*-4-(3,4-difluorophenyl)-3-(1-(4-(2-cyanophenyl)cyclohexyl)-3-hydroxy-azetidin-
3-yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2-one;

trans-4-(3,4-difluorophenyl)-3-(1-(4-(2-cyanophenyl)cyclohexyl)azetidin-3-
yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2-one;


25

trans-4-(3,4-difluorophenyl)-3-(1-(4-(2-pyridyl)cyclohexyl)azetidin-3-
yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2(3H)-one;

and pharmaceutically acceptable salts thereof.

30

In a third class of the invention is a compound of Formula (V):



wherein

- 5 each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

- 10 each X³ is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, C₁-₄ alkyl, or CF₃;

R² is hydrogen or C₁-C₄ alkyl;

- 15 R³ is hydrogen or C₁-C₄ alkyl;

R⁹ is hydrogen or C₁-C₄ alkyl;

- 20 R¹⁰ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

- 25 R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

t is an integer from 2 to 4; and

5

q₁, q₂ and q₃ are each independently integers from 0 to 4;

or a pharmaceutically acceptable salt thereof.

10

In a sub-class of the third class is a compound of formula (V), wherein

each X¹ is independently hydrogen, halogen, cyano, methyl, ethyl, CF₃, or OCF₃;

each X² is hydrogen;

15

each X³ is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy, ethoxy, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, CF₃, or OCF₃;

R² is hydrogen;

20

R³ is hydrogen;

R⁹ is hydrogen;

25

R¹⁰ is hydrogen, cyano, or OR^c;

R^c is hydrogen, methyl, or ethyl;

t is 2 or 3;


30

q₁ and q₃ are each independently integers from 0 to 3; and

all other variables are as defined in the third class;

or a pharmaceutically acceptable salt thereof.

In a fourth class of the invention is a compound of Formula (VI):



5

wherein

each X^1 is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

10

each X^2 is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

each X^3 is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂,
15 SO₂N(R^c)₂, SO₂R^c, C₁-C₄ alkyl, or CF₃;

R^2 is hydrogen or C₁-C₄ alkyl;

R^3 is hydrogen or C₁-C₄ alkyl;

20

R^9 is hydrogen or C₁-C₄ alkyl;

R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

25 R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

t is an integer from 2 to 4; and

q₁, q₂ and q₃ are each independently integers from 0 to 4;

5 or a pharmaceutically acceptable salt thereof.

In a sub-class of the fourth class is a compound of formula (VI),
wherein

10 each X¹ is independently hydrogen, halogen, cyano, methyl, ethyl, CF₃, or OCF₃;

each X² is hydrogen;

each X³ is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy,
15 ethoxy, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, CF₃, or OCF₃;

R² is hydrogen;

R³ is hydrogen;

20

R⁹ is hydrogen;

R^c is hydrogen, methyl, or ethyl;

25 t is 2 or 3;


q₁ and q₃ are each independently integers from 0 to 3; and

all other variables are as defined in the fourth class;

30

or a pharmaceutically acceptable salt thereof.

In a fifth class of the invention is a compound of Formula (VII):



wherein

5 each X^1 is independently hydrogen, halogen, cyano, C_1 - C_4 alkyl, CF_3 , or $(CH_2)_0$ - $4OR^a$;

each X^2 is independently hydrogen, halogen, cyano, C_1 - C_4 alkyl, CF_3 , or $(CH_2)_0$ - $4OR^a$;

10 each X^3 is independently hydrogen, halogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, $SO_2N(R^c)_2$, SO_2R^c , C_1 - C_4 alkyl, or CF_3 ;

R^2 is hydrogen or C_1 - C_4 alkyl;

15 R^3 is hydrogen or C_1 - C_4 alkyl;

R^a is hydrogen, C_1 - C_4 alkyl, or $(CH_2)_0$ - $3CF_3$;

20 R^c is hydrogen, C_1 - C_4 alkyl, or $(CH_2)_0$ - $3CF_3$;

o and p are each independently 0 or 1;

w is an integer from 0 to 2; and

25 q_1 , q_2 and q_3 are each independently integers from 0 to 4;

or a pharmaceutically acceptable salt thereof.

In a sub-class of the fifth class is a compound of formula (VII),
wherein

5

each X^1 is independently hydrogen, halogen, cyano, methyl, ethyl, CF_3 , or OCF_3 ;

each X^2 is hydrogen;

10 each X^3 is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy, ethoxy, CO_2R^c , $CON(R^c)_2$, $SO_2N(R^c)_2$, SO_2R^c , CF_3 , or OCF_3 ;

R^2 is hydrogen;

15 R^3 is hydrogen;

R^c is hydrogen, methyl, or ethyl;

w is 0 or 1;


20

q_1 and q_3 are each independently integers from 0 to 3; and

all other variables are as defined in the fifth class;

25 or a pharmaceutically acceptable salt thereof.

In a sixth class of the invention is a compound of Formula (VIII):



wherein

5 each X^1 is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

each X^2 is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

10 each X^3 is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, C₁-C₄ alkyl, or CF₃;

R² is hydrogen or C₁-C₄ alkyl;

15 R⁹ is hydrogen or C₁-C₄ alkyl;

R¹⁰ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the phenyl are independently selected
 20 from halogen, cyano, OR^c, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

25 R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

o and p are each independently 0 or 1;

w is an integer from 0 to 2;

5

q₁, q₂ and q₃ are each independently integers from 0 to 4;

or a pharmaceutically acceptable salt thereof.

10

In a sub-class of the sixth class is a compound of formula (VIII),

wherein

each X¹ is independently hydrogen, halogen, cyano, methyl, ethyl, CF₃, or OCF₃;

15

each X² is hydrogen;

each X³ is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy, ethoxy, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, CF₃, or OCF₃;

20

R² is hydrogen;

R⁹ is hydrogen;

R¹⁰ is hydrogen, cyano, or OR^c;

25

R^c is hydrogen, methyl, or ethyl;

w is 0 or 1;


30

q₁ and q₃ are each independently integers from 0 to 3; and

all other variables are as defined in the sixth class;

or a pharmaceutically acceptable salt thereof.

In a seventh class of the invention is a compound of Formula (IX):



5 wherein

each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

10 each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

each X³ is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, C₁₋₄ alkyl, or CF₃;

15 R² is hydrogen or C₁-C₄ alkyl;

R⁹ is hydrogen or C₁-C₄ alkyl;

20 R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

o and p are each independently 0 or 1;

25 w is an integer from 0 to 2; and

q1, q2 and q3 are each independently integers from 0 to 4;

or a pharmaceutically acceptable salt thereof.

5

In a sub-class of the seventh class is a compound of formula (IX),
wherein

each X^1 is independently hydrogen, halogen, cyano, methyl, ethyl, CF_3 , or OCF_3 ;

10

each X^2 is hydrogen;

each X^3 is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy,
ethoxy, CO_2R^c , $CON(R^c)_2$, $SO_2N(R^c)_2$, SO_2R^c , CF_3 , or OCF_3 ;

15

R^2 is hydrogen;

R^9 is hydrogen;

20

R^c is hydrogen, methyl, or ethyl;

w is 0 or 1;

q1 and q3 are each independently integers from 0 to 3; and

25

all other variables are as defined in the seventh class;

or a pharmaceutically acceptable salt thereof.

30

The present invention also includes a pharmaceutical composition comprising a therapeutically effective amount of any of the compounds described above and a pharmaceutically acceptable carrier. In one embodiment is a pharmaceutical composition made by combining any of the compounds described above and a pharmaceutically acceptable carrier. The present invention further

includes a process for making a pharmaceutical composition comprising combining any of the compounds described above and a pharmaceutically acceptable carrier.

The present invention further includes a pharmaceutical composition as described in the preceding paragraph further comprising a therapeutically effective amount of a testosterone 5-alpha reductase inhibitor. In one embodiment, the testosterone 5-alpha reductase inhibitor is a type 1, a type 2, both a type 1 and a type 2 (i.e., a three component combination comprising any of the compounds described above combined with both a type 1 testosterone 5-alpha reductase inhibitor and a type 2 testosterone 5-alpha reductase inhibitor), or a dual type 1 and type 2 testosterone 5-alpha reductase inhibitor. In another embodiment, the testosterone 5-alpha reductase inhibitor is a type 2 testosterone 5-alpha reductase inhibitor. The testosterone 5-alpha reductase inhibitor is suitably finasteride.

The present invention also includes a method of treating benign prostatic hyperplasia in a subject in need thereof which comprises administering to the subject a therapeutically effective amount of any of the compounds (or any of the compositions) described above. In one embodiment of the method of treating BPH, the compound (or composition) does not cause a fall in blood pressure at dosages effective to alleviate BPH. In another embodiment of the method of treating BPH, the compound is administered in combination with a testosterone 5-alpha reductase inhibitor. A suitable testosterone 5-alpha reductase inhibitor for use in the method is finasteride.

The present invention also includes a method of inhibiting contraction of prostate tissue or relaxing lower urinary tract tissue in a subject in need thereof which comprises administering to the subject a therapeutically effective amount of any of the compounds (or any of the compositions) described above. In one embodiment of the method of inhibiting contraction of prostate tissue or relaxing lower urinary tract tissue, the compound (or composition) additionally does not cause a fall in blood pressure at dosages effective to inhibit contraction of prostate tissue. In another embodiment, the compound is administered in combination with a testosterone 5-alpha reductase inhibitor; the testosterone 5-alpha reductase inhibitor is suitably finasteride.

The present invention also includes a method of treating a disease which is susceptible to treatment by antagonism of the alpha 1a receptor which comprises administering to a subject in need thereof an amount of any of the compounds described above effective to treat the disease. Diseases which are

susceptible to treatment by antagonism of the alpha 1a receptor include, but are not limited to, BPH, high intraocular pressure, high cholesterol, impotency, sympathetically mediated pain, migraine (see Vatz, *Headache* (1997), 37: 107-108) and cardiac arrhythmia.

- 5 The present invention also includes the use of any of the compounds described above in the preparation of a medicament for: a) treating benign prostatic hyperplasia; b) relaxing lower urinary tract tissue; or c) inhibiting contraction of prostate tissue; in a subject in need thereof.

- The present invention further includes the use of any of the alpha 1a
10 antagonist compounds described above and a 5-alpha reductase inhibitor for the manufacture of a medicament for: a) treating benign prostatic hyperplasia; b) relaxing lower urinary tract tissue; or c) inhibiting contraction of prostate tissue which comprises an effective amount of the alpha 1a antagonist compound and an effective amount of 5-alpha reductase inhibitor, together or separately.

- 15 As used herein, the term "C₁-C₈ alkyl" means linear or branched chain alkyl groups having from 1 to 8 carbon atoms and includes all of the octyl alkyl, heptyl alkyl, hexyl alkyl and pentyl alkyl isomers as well as n-, iso-, sec- and t-butyl, n- and isopropyl, ethyl and methyl. "C₁-C₄ alkyl" means n-, iso-, sec- and t-butyl, n- and isopropyl, ethyl and methyl.

- 20 The term "C₁-C₆ alkoxy" means an -O-alkyl group wherein alkyl is C₁ to C₆ alkyl. "C₁-C₄ alkoxy" has an analogous meaning; i.e., it is an alkoxy group selected from methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, tert-butoxy, and sec-butoxy.

- The term "C₃-C₈ cycloalkyl" means a cyclic ring of an alkane having
25 three to eight total carbon atoms (i.e., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl). "C₃-C₆ cycloalkyl" refers to a cyclic ring selected from cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

- The term "C₁-C₈ alkyl- C₃-C₈ cycloalkyl" means a C₃-C₈ cycloalkyl
group as defined above substituted with one or more linear or branched chain alkyl
30 groups having from 1 to 8 carbon atoms.

 The term "C₃-C₈ cycloalkyl- C₁-C₈ alkyl" means a C₁-C₈ alkyl group as defined above substituted with one or more cyclic alkyl groups having from 3 to 8 carbon atoms.

The term "halogen" (which may alternatively be referred to as "halo") refers to fluorine, chlorine, bromine and iodine (alternatively, fluoro, chloro, bromo, and iodo).

The term "fluorinated C₁-C₈ alkyl" (which may alternatively be referred to as "C₁-C₈ fluoroalkyl") means a C₁ to C₈ linear or branched alkyl group as defined above with one or more fluorine substituents. The term "fluorinated C₁-C₄ alkyl" has an analogous meaning. Representative examples of suitable fluoroalkyls include the series (CH₂)₀₋₄CF₃ (i.e., trifluoromethyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoro-n-propyl, etc.), 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 3,3,3-trifluoroisopropyl, 1,1,1,3,3,3-hexafluoroisopropyl, and perfluorohexyl.

The term "fluorinated C₃-C₈ cycloalkyl" (which may alternatively be referred to as "C₃-C₈ fluorocycloalkyl") means a cycloalkyl group as defined above with one or more fluorine substituents. "Fluorinated C₃-C₆ cycloalkyl" has an analogous meaning. Representative examples of suitable fluorocycloalkyls include all isomers of fluorocyclohexyl (i.e., 1-, 2-, 3-, and 4-fluorocyclohexyl), difluorocyclohexyl (e.g., 2,4-difluorocyclohexyl, 3,4-difluorocyclohexyl, etc.), fluorocyclopentyl, and so forth.

The term "fluorinated C₁-C₆ alkoxy" (which may alternatively be referred to as "C₁-C₆ fluoroalkoxy") means a C₁-C₆ alkoxy group as defined above wherein the alkyl moiety has one or more fluorine substituents. "Fluorinated C₁-C₄ alkoxy" has an analogous meaning. Representative examples include the series O(CH₂)₀₋₄CF₃ (i.e., trifluoromethoxy, 2,2,2-trifluoroethoxy, 3,3,3-trifluoro-n-propoxy, etc.), 1,1,1,3,3,3-hexafluoroisopropoxy, and so forth.

The term "heterocyclic" (which may alternatively be referred to as "heterocycle") refers to a stable 5- to 7-membered monocyclic ring system which may be saturated or unsaturated; which consists of carbon atoms and from one to three heteroatoms selected from N, O or S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized; and which is optionally substituted with one or more substituent groups including, but not limited to, halo, cyano, alkyl, fluoroalkyl, alkoxy, fluoroalkoxy, alkylalkoxy, cycloalkyl, fluorocycloalkyl, amino, aryl (e.g., phenyl), carboxy, carboxylate, sulfonamido, sulfonyl, and the like. Suitable heterocyclic groups include, but are not limited to, piperidinyl, piperazinyl, oxopiperazinyl, oxopiperidinyl, oxopyrrolidinyl, oxoazepinyl, azepinyl, pyrrolyl, pyrrolidinyl, furanyl, thienyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, pyrazinyl,

pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isooxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, thiadiazolyl, tetrahydropyranlyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl. Morpholino is the same as morpholinyl.

- 5 The term "thienyl," as used herein, refers to the group



The term "aryl" refers herein to aromatic mono- and poly-carbocyclic ring systems, optionally substituted, including, but not limited to, phenyl, substituted phenyl, naphthyl, and substituted naphthyl.

- 10 The term "heteroaryl" refers to aromatic heterocyclic or substituted aromatic heterocyclic ring systems, including, but not limited to, pyridyl, substituted pyridyl, pyridyl N-oxide ($N\text{O}$), substituted pyridyl N-oxide, pyrazinyl, substituted pyrazinyl, thienyl, substituted thienyl, thiazolyl, substituted thiazolyl, furanyl, substituted furanyl, quinazolinyl, and substituted quinazolinyl.

- 15 The term "substituted" includes mono- and poly-substitution by a named substituent to the extent such single and multiple substitution is chemically allowed.

- It is understood that the definition of a substituent (e.g., CO_2R^c) or variable (e.g., R^c) at a particular location in a molecule is independent of its definitions at other locations in that molecule. Thus, for example, when R^4 is mono-substituted phenyl wherein the substituent is $\text{CO}_2\text{R}^c = \text{CO}_2\text{H}$, and R^5 is also mono-substituted phenyl wherein the substituent is CO_2R^c , it is understood that the substituent on the phenyl in R^5 can be any one of CO_2H , CO_2Me , CO_2Et , CO_2Pr , $\text{CH}_2\text{CO}_2\text{H}$, $\text{CH}_2\text{CO}_2\text{Me}$, $\text{CH}_2\text{CO}_2\text{Et}$, $\text{CH}_2\text{CO}_2\text{Pr}$, $(\text{CH}_2)_2\text{CO}_2\text{H}$, etc. As another example, $\text{NRC}(\text{=O})\text{R}^c$ represents $\text{NHC}(\text{=O})\text{H}$, $\text{NHC}(\text{=O})\text{Me}$, $\text{NMeC}(\text{=O})\text{Me}$, $\text{NMeC}(\text{=O})\text{Et}$, etc.
- 25

- It is also understood that the definition of a substituent or variable at a particular location in a molecule is independent of the definition of another occurrence of the same substituent or variable at the same location. Thus, $\text{C}(\text{=O})\text{N}(\text{R}^c)_2$ represents groups such as $-\text{C}(\text{=O})\text{NH}_2$, $-\text{C}(\text{=O})\text{NHCH}_3$, $-\text{C}(\text{=O})\text{NHC}_2\text{H}_5$, $-\text{C}(\text{=O})\text{N}(\text{CH}_3)\text{C}_2\text{H}_5$, etc.
- 30

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by the methods set forth below and, when viewed in the light of this disclosure, by techniques known in the art. Where multiple substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally.

Representative embodiments for the variables and substituents set forth in Formula (I) include the following:


10 A^1 , A^2 , A^3 and A^4 are each independently selected from $C-X^2$ or N, provided that no more than two of A^1 , A^2 , A^3 and A^4 are nitrogen, and further provided that when two of A^1 , A^2 , A^3 and A^4 are N, each nitrogen atom is bonded to two carbon atoms. In one embodiment, A^1 and A^4 are nitrogen and A^2 and A^3 are $C-X^2$. In another embodiment, A^2 and A^4 are nitrogen and A^1 and A^3 are $C-X^2$. In yet another embodiment, one of A^1 , A^2 , A^3 and A^4 is nitrogen, and the others are $C-X^2$. In still another embodiment, all of A^1 , A^2 , A^3 and A^4 are $C-X^2$.

Each X^1 is independently hydrogen, halogen, cyano, nitro, C_1 -C8 alkyl, C_3 -C8 cycloalkyl, fluorinated C_1 -C8 alkyl, fluorinated C_3 -C8 cycloalkyl, or $(CH_2)_0-4OR^a$; or is independently hydrogen, halogen, cyano, C_1 -C4 alkyl, fluorinated C_1 -C4 alkyl, C_3 -C6 cycloalkyl, fluorinated C_3 -C6 cycloalkyl, or $(CH_2)_0-4OR^a$; or is independently hydrogen, halogen, cyano, C_1 -C4 alkyl, $(CH_2)_0-4OCF_3$, or $(CH_2)_0-4CF_3$; or is independently hydrogen, halogen, cyano, C_1 -C4 alkyl, CF_3 , or $(CH_2)_0-4OR^a$. In another embodiment, each X^1 is independently hydrogen, halogen, cyano, methyl, ethyl, CF_3 , or OCF_3 . In another embodiment, each X^1 is independently hydrogen, fluorine, cyano, methyl, ethyl, CF_3 , or OCF_3 . In still another embodiment, each X^1 is fluorine.

Each X^2 is independently hydrogen, halogen, cyano, nitro, C_1 -C8 alkyl, C_3 -C8 cycloalkyl, fluorinated C_1 -C8 alkyl, fluorinated C_3 -C8 cycloalkyl, or $(CH_2)_0-4OR^a$; or is independently hydrogen, halogen, cyano, C_1 -C4 alkyl, fluorinated C_1 -C4 alkyl, C_3 -C6 cycloalkyl, fluorinated C_3 -C6 cycloalkyl, or $(CH_2)_0-4OR^a$; or is independently hydrogen, halogen, cyano, C_1 -C4 alkyl, $(CH_2)_0-4OCF_3$, or $(CH_2)_0-4CF_3$; or is independently hydrogen, halogen, cyano, C_1 -C4 alkyl, CF_3 , or $(CH_2)_0-4OR^a$. In another embodiment, each X^2 is independently hydrogen, halogen,

cyano, methyl, ethyl, CF₃, or OCF₃. In still another embodiment, each X² is hydrogen.

One aspect of the embodiment in which all of A¹, A², A³ and A⁴ are C-X² is the dihydroquinazolin-2-one moiety of formula:



5

wherein X² is as defined above; and q₂ is an integer from 0 to 4; or from 0 to 3; or from 0 to 2; or is 0 or 1; or is 0; or is 1.

R¹ is hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl, phenyl, or substituted phenyl; or is hydrogen, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, phenyl, or substituted phenyl; or is hydrogen, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, phenyl, or mono- or di- or tri-substituted phenyl. In another embodiment, R¹ is hydrogen.

In R¹, the substituents on the substituted phenyl are independently selected from halogen, cyano, nitro, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl and (CH₂)₀₋₄OR^a; or are independently selected from halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl and (CH₂)₀₋₄OR^a.

R² is hydrogen, C₁-C₈ alkyl, or fluorinated C₁-C₈ alkyl; or is hydrogen, C₁-C₄ alkyl, or fluorinated C₁-C₄ alkyl; or is hydrogen or C₁-C₄ alkyl. In another embodiment, R² is hydrogen.

R³ is hydrogen, C₁-C₈ alkyl, or fluorinated C₁-C₈ alkyl; or is hydrogen, C₁-C₄ alkyl, or fluorinated C₁-C₄ alkyl; or is hydrogen or C₁-C₄ alkyl. In another embodiment, R³ is hydrogen.

Y is carbon or nitrogen, provided that when Y is nitrogen, R⁵ is absent. In one embodiment, Y is carbon.

Z is CH₂, CHOR^b, CHF, CHR^b, C(R^b)₂, CF₂, CHCHF₂, C=CF₂, or C=O. In one embodiment, Z is CH₂, CHF, CHR^b, C(R^b)₂, CF₂, CHCHF₂, or

C=CF₂. In another embodiment, Z is CH₂, CHOH, or C=O, provided that when u is zero, Z is CH₂.

E, G, L and M are each independently selected from hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, (CH₂)₀₋₄OR^c, (CH₂)₀₋₄N(R^c)₂, (CH₂)₀₋₄CN, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂R^c and (CH₂)₀₋₄SO₂N(R^c)₂; or are each independently selected from hydrogen, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; or are each independently selected from hydrogen, C₁-C₄ alkyl and (CH₂)₀₋₃CF₃; or are each independently selected from hydrogen and C₁-C₄ alkyl. In one embodiment, E, G, L and M are all hydrogen.


J is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, (CH₂)₁₋₄OR^c, (CH₂)₁₋₄N(R^c)₂, (CH₂)₁₋₄CN, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂R^c, or (CH₂)₀₋₄SO₂N(R^c)₂; or is hydrogen, C₁-C₄ alkyl or fluorinated C₁-C₄ alkyl; or is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃; or is hydrogen or C₁-C₄ alkyl; or is hydrogen.

R⁴ is phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, thienyl, furanyl, substituted pyridyl, substituted thienyl, or substituted furanyl; or is phenyl, mono- or di- or tri-substituted phenyl, pyridyl, thienyl, or furanyl; or is phenyl or mono- or di- or tri-substituted phenyl; or is phenyl or mono-substituted phenyl.

In R⁴, the substituents on the substituted phenyl are independently selected from halogen, cyano, OR^c, (CH₂)₀₋₃CON(R^c)₂, (CH₂)₀₋₃CO₂R^c, methylenedioxy when the phenyl ring is di-substituted and the substituents are on adjacent carbon atoms, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; or are independently selected from halogen, cyano, CO₂R^c, OR^c, CON(R^c)₂, methylenedioxy, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; or are independently selected from CF₃, cyano, halogen (e.g., fluorine), and C₁-C₄ alkyl.

In R⁴, the substituents on the substituted naphthyl, pyridyl, thienyl, or furanyl are independently selected from phenyl, OR^c, halogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl, C₃-C₈ cycloalkyl and fluorinated C₃-C₈ cycloalkyl; or are independently selected from CF₃, phenyl, OR^c, halogen, and C₁-C₄ alkyl.

In one embodiment, R⁴ is represented by the formula:



wherein each X^3 is independently halogen, cyano, OR^c , $(CH_2)_{0-3}CON(R^c)_2$, $(CH_2)_{0-3}CO_2R^c$, C_1 - C_4 alkyl or fluorinated C_1 - C_4 alkyl; and q_3 is an integer of from 0 to 5; or from 0 to 4; or from 0 to 3; or from 0 to 2; or is 0 or 1. In other

- 5 embodiments, each X^3 is independently hydrogen, halogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, $SO_2N(R^c)_2$, SO_2R^c , C_1 - C_4 alkyl, or CF_3 ; or is independently hydrogen, CF_3 , cyano, halogen, or C_1 - C_4 alkyl; or is independently hydrogen, CF_3 , cyano, fluorine, methyl, or ethyl; or is independently hydrogen, CF_3 , cyano, or fluorine.

- R^5 is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, tetrazole, 10 isooxadiazole, phenyl, or substituted phenyl; or is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, phenyl, or mono- or di- or tri-substituted phenyl; or is hydrogen, cyano, CO_2R^c , $CON(R^c)_2$, tetrazole and isooxadiazole; or is hydrogen, cyano, or CO_2R^c ; or is cyano or CO_2R^c . In one embodiment, R^5 is cyano or CO_2CH_3 .

- In R^5 , the substituents on the substituted phenyl are independently 15 selected from halogen, cyano, $(CH_2)_{0-3}CON(R^c)_2$, $(CH_2)_{0-3}CO_2R^c$, methylenedioxy when the phenyl ring is di-substituted and the substituents are on adjacent carbon atoms, C_1 - C_4 alkyl and fluorinated C_1 - C_4 alkyl; or are independently selected from halogen, cyano, CO_2R^c , OR^c , $CON(R^c)_2$, methylenedioxy, C_1 - C_4 alkyl and fluorinated C_1 - C_4 alkyl.

- 20 R^6 and R^7 are each independently selected from hydrogen, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, fluorinated C_1 - C_8 alkyl and fluorinated C_3 - C_8 cycloalkyl; or one of R^6 and R^7 is hydrogen, and the other of R^6 and R^7 is hydrogen or C_1 - C_4 alkyl. In another embodiment, R^6 and R^7 are both hydrogen. In still another embodiment, R^6 and R^7 together with the carbon atom to which they are attached 25 form methylene ($-CH_2-$) units or alkylidene units of formula $(-CR^wH-)$ wherein R^w is C_1 - C_4 alkyl or, in cases where there are at least two CR^6R^7 's (e.g., when $w = 2$), mixtures of the foregoing units.


- R^8 is phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, substituted pyridyl, pyridyl N-oxide ($N\text{O}$), substituted pyridyl N-oxide, 30 pyrazinyl, substituted pyrazinyl, thienyl, substituted thienyl, thiazolyl, substituted thiazolyl, furanyl, substituted furanyl, quinazolinyl, or substituted quinazolinyl; or is

phenyl, mono- or di- or tri-substituted phenyl, pyridyl, or mono- or di- or tri-substituted pyridyl; or is phenyl or mono- or di- or tri-substituted phenyl.

- In R^8 , the substituents on the phenyl are independently selected from halogen, cyano, nitro, OR^c , $N(R^c)_2$, NR^cCOR^c , $NR^cCON(R^c)_2$, $NR^cSO_2R^c$, $NR^cSO_2N(R^c)_2$, $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$, C₁₋₄ alkyl and fluorinated C₁₋₄ alkyl; or are independently selected from halogen, cyano, OR^c , $N(R^c)_2$, NR^cCOR^c , $NR^cCON(R^c)_2$, $NR^cSO_2R^c$, $NR^cSO_2N(R^c)_2$, $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$, C₁₋₄ alkyl and CF_3 ; or are independently selected from CF_3 , cyano, halogen (e.g., fluorine), and C₁₋₄ alkyl.

- In R^8 , the substituents on the substituted naphthyl, pyridyl, pyrazinyl, thienyl, thiazolyl, furanyl, or quinazolinyl are independently selected from cyano, nitro, $N(R^c)_2$, $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$, phenyl, OR^c , halogen, C₁₋₄ alkyl, fluorinated C₁₋₄ alkyl, C₃₋₈ cycloalkyl and fluorinated C₃₋₈ cycloalkyl; or are independently selected from cyano, nitro, $N(R^c)_2$, $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$, phenyl, OR^c , halogen, C₁₋₄ alkyl and CF_3 .

In one embodiment, R^8 is represented by the formula:



wherein X^3 and q_3 are as defined above.


- R^9 is hydrogen, C₁₋₈ alkyl, or fluorinated C₁₋₈ alkyl; or is hydrogen, C₁₋₄ alkyl, or fluorinated C₁₋₄ alkyl. In another embodiment, R^2 is hydrogen.

- R^{10} is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, tetrazole, isooxadiazole, phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, substituted pyridyl, thienyl, substituted thienyl, furanyl, or substituted furanyl; or is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, phenyl, or mono- or di- or tri-substituted phenyl; or is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, phenyl, or mono- or di-substituted phenyl; or is hydrogen, cyano, or OR^c .

In R^{10} , the substituents on the substituted phenyl are independently selected from halogen, cyano, nitro, OR^c , $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $N(R^c)_2$, NR^cCOR^c , $NR^cCON(R^c)_2$, $NR^cSO_2R^c$, $NR^cSO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; or are
 5 independently selected from halogen, cyano, OR^c , $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $N(R^c)_2$, NR^cCOR^c , $NR^cCON(R^c)_2$, $NR^cSO_2R^c$, $NR^cSO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$ and C₁-C₄ alkyl; or are independently selected from halogen, cyano, OR^c , $(CH_2)_{0-2}CO_2R^c$, $(CH_2)_{0-2}CON(R^c)_2$ and C₁-C₄ alkyl.

10 In R^{10} , the substituents on the substituted naphthyl, pyridyl, thienyl, or furanyl are independently selected from $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$, phenyl, OR^c , halogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl and C₃-C₈ cycloalkyl; or are independently selected from CF₃, $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$,
 15 phenyl, OR^c , halogen, C₁-C₄ alkyl and C₃-C₈ cycloalkyl.

In one embodiment, R^{10} is represented by the formula:



wherein X^3 and q_3 are as defined above.

20 R^{13} , R^{14} , R^{15} and R^{16} are each independently selected from hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, and $(CH_2)_{2-4}OR^c$; or are each independently selected from hydrogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl, and $(CH_2)_{2-4}OR^b$; or are each independently selected from hydrogen, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; or are each
 25 independently selected from hydrogen, C₁-C₄ alkyl and $(CH_2)_{1-4}CF_3$; or are each independently selected from hydrogen and C₁-C₄ alkyl. In one embodiment, R^{13} , R^{14} , R^{15} and R^{16} are all hydrogen.

R^{18} and R^{20} are each independently selected from hydrogen and OR^d ; or are each independently selected from hydrogen, hydroxy, methoxy, ethoxy, and
 30 CF_3 ; or are each independently selected from hydrogen and hydroxy (e.g., R^{18} and R^{20} are both hydrogen or both hydroxy).

R^a is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, or C₃-C₈ cycloalkyl; or is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃; or is hydrogen, methyl, ethyl, or CF₃; or is hydrogen, methyl, or ethyl.

R^b is C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or
5 fluorinated C₃-C₈ cycloalkyl; or is C₁-C₄ alkyl or (CH₂)₀₋₃CF₃; or is methyl, ethyl, or CF₃; or is methyl or ethyl.

R^c is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or fluorinated C₃-C₈ cycloalkyl; or is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃; or is hydrogen, methyl, ethyl, or CF₃; or is hydrogen, methyl, or ethyl.

10 R^d is hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or (CH₂)₀₋₄CF₃; or is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃; or is hydrogen, methyl, ethyl, or CF₃; or is hydrogen, methyl, or ethyl.

m, n, o, and p are each independently integers from 0 to 3; or from 0 to 2; or are 0 or 1. In other embodiments, m and n are each integers of from 0 to 3,
15 wherein the sum of m + n is an integer of from 0 to 3; or m and n are each integers of from 0 to 2, wherein the sum of m + n is an integer of from 0 to 2; or m and n are each either 1 or 2 and the sum of m + n is 1 or 2 (e.g., m = 1 and n = 1). In still other embodiments, o and p are each integers of from 0 to 3, wherein the sum of o + p is an integer of from 0 to 3; or o and p are each integers of from 0 to 2, wherein the sum of
20 o + p is an integer of from 0 to 2; or o and p are each either 1 or 2, and the sum of o + p is either 1 or 2 (e.g., o = 1 and p = 1).

q₁ is an integer from 0 to 5; or from 0 to 4; or from 0 to 3; or from 0 to 2; or is 0 or 1. In one embodiment, q₁ is 2. In one aspect of the embodiment of q₁ = 2, the substituted phenyl is either 2,4- or 3,4-di-substituted (e.g., 3,4-difluorophenyl).

25 t is an integer from 2 to 5; or from 2 to 4; or is 2 or 3; or is 2; or is 3.

u and v are each independently integers from 0 to 3; provided that u and v are not both zero; and further provided that when u is zero, Z is selected from CH₂, CHF, CHR^b, C(R^b)₂, CF₂, CHCHF₂ and C=CF₂.

w is an integer from 0 to 3, provided that when w is 0, R²⁰ is
30 hydrogen. In other embodiments w is an integer from 0 to 2; or is 0 or 1; or is 0; or is 1.

Representative compounds of the present invention exhibit selectivity for the human alpha 1a adrenergic receptor. One implication of this selectivity is that

these compounds display selectivity for lowering intraurethral pressure without substantially affecting diastolic blood pressure.

Representative compounds of this invention display submicromolar affinity for the human alpha 1a adrenergic receptor subtype while displaying lower
5 affinity for the human alpha 1d and alpha 1b adrenergic receptor subtypes, and many other G-protein coupled human receptors. Particular representative compounds of this invention exhibit nanomolar and subnanomolar affinity for the human alpha 1a adrenergic receptor subtype while displaying at least about 10 fold lower affinity for the human alpha 1d and alpha 1b adrenergic receptor subtypes, and many other G-
10 protein coupled human receptors (e.g., serotonin, dopamine, alpha 2 adrenergic, beta adrenergic or muscarinic receptors). Still other compounds of this invention exhibit nanomolar and subnanomolar affinity for the human alpha 1a adrenergic receptor subtype while displaying at least about 20 fold lower affinity for the human alpha 1d and alpha 1b adrenergic receptor subtypes, in addition to exhibiting selectivity over
15 other G-protein coupled human receptors (e.g., serotonin, dopamine, alpha 2 adrenergic, beta adrenergic or muscarinic receptors).

These compounds are administered in dosages effective to antagonize the alpha 1a receptor where such treatment is needed; e.g., treatment of BPH. For use in medicine, the salts of the compounds of this invention refer to non-toxic
20 "pharmaceutically acceptable salts." Other salts may, however, be useful in the preparation of the compounds according to the invention or in the preparation of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a
25 solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth
30 metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium salts. Thus, representative pharmaceutically acceptable salts include the following:

Acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, camsylate, carbonate, clavulanate, citrate, dihydrochloride, edetate, edisylate,
35 estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate,

- hexylresorcinate hydrobromide, hydrochloride, hydroxynaphthoate, hydroiodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, n-methylglucamine ammonium salt, oleate, palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, tosylate and valerate.

Compounds of this invention are used to reduce the acute symptoms of BPH. Thus, compounds of this invention may be used alone or in combination with more long-term anti-BPH therapeutics, such as testosterone 5- α reductase inhibitors, including PROSCAR® (finasteride). Aside from their utility as anti-BPH agents, these compounds may be used to induce highly tissue-specific, localized α 1a adrenergic receptor blockade whenever this is desired. Effects of this blockade include reduction of intra-ocular pressure, control of cardiac arrhythmias, and possibly a host of α 1a receptor mediated central nervous system events.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds of this invention which are readily convertible in vivo into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various conditions described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound in vivo after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in Design of Prodrugs, ed. H. Bundgaard, Elsevier, 1985.

The present invention further includes metabolites of the compounds of the present invention. Metabolites include active species produced upon introduction of compounds of this invention into the biological milieu.

Where the compounds according to the invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds according to the invention possess two or more chiral centers, they may additionally exist as diastereoisomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for compounds of the present invention may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds of the present invention may form solvates with

water (i.e., hydrates) or common organic solvents. Such solvates are also encompassed within the scope of this invention.

The term "selective alpha 1a adrenergic receptor antagonist," as used herein, refers to an alpha 1a antagonist compound which is at least about ten fold
5 selective for the human alpha 1a adrenergic receptor as compared to the human alpha 1b, alpha 1d, alpha 2a, alpha 2b and alpha 2c adrenergic receptors.

The term "lower urinary tract tissue," as used herein, refers to and includes, but is not limited to, prostatic smooth muscle, the prostatic capsule, the urethra and the bladder neck.

10 The term "subject," as used herein refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

The term "therapeutically effective amount" as used herein means that amount of active compound or pharmaceutical agent that elicits the biological
15 or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease being treated.

The present invention includes pharmaceutical compositions comprising one or more compounds of this invention in association with a
20 pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the compositions may
25 be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch,
30 lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is
35 dispersed evenly throughout the composition so that the composition may be readily

subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated
5 or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the
10 duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

As used herein, the term "composition" encompasses a product
15 comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include
20 aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose,
25 polyvinyl-pyrrolidone or gelatin.

Where the processes for the preparation of the compounds according to the invention give rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds
30 may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also
35 be resolved by formation of diastereomeric esters or amides, followed by

chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known in the art.

The specificity of binding of compounds showing affinity for the alpha 1a receptor is shown by comparing affinity to membranes obtained from transfected cell lines that express the alpha 1a receptor and membranes from cell lines or tissues known to express other types of alpha (e.g., alpha 1d, alpha 1b) or beta adrenergic receptors. Expression of the cloned human alpha 1d, alpha 1b, and alpha 1a receptors and comparison of their binding properties with known selective antagonists provides a rational way for selection of compounds and discovery of new compounds with predictable pharmacological activities. Antagonism by these compounds of the human alpha 1a adrenergic receptor subtype may be functionally demonstrated in anesthetized animals. These compounds may be used to increase urine flow without exhibiting hypotensive effects.

The ability of compounds of the present invention to specifically bind to the alpha 1a receptor makes them useful for the treatment of BPH. The specificity of binding of compounds showing affinity for the alpha 1a receptor is compared against the binding affinities to other types of alpha or beta adrenergic receptors. The human alpha adrenergic receptor of the 1a subtype was recently identified, cloned and expressed as described in PCT International Application Publication Nos. WO94/08040, published 14 April 1994 and WO 94/21660, published 29 September 1994. The cloned human alpha 1a receptor, when expressed in mammalian cell lines, is used to discover ligands that bind to the receptor and alter its function. Expression of the cloned human alpha 1d, alpha 1b, and alpha 1a receptors and comparison of their binding properties with known selective antagonists provides a rational way for selection of compounds and discovery of new compounds with predictable pharmacological activities.

Compounds of this invention exhibiting human alpha 1a adrenergic receptor antagonism may further be defined by counterscreening. This is accomplished according to methods known in the art using other receptors responsible for mediating diverse biological functions. [See e.g., PCT International Application Publication No. WO94/10989, published 26 May 1994; US 5403847, issued April 4, 1995]. Compounds which are both selective amongst the various human alpha 1a adrenergic receptor subtypes and which have low affinity for other receptors, such as the alpha 2 adrenergic receptors, the β -adrenergic receptors, the muscarinic receptors, the serotonin receptors, the histamine receptors, and others are particularly preferred.

The absence of these non-specific activities may be confirmed by using cloned and expressed receptors in an analogous fashion to the method disclosed herein for identifying compounds which have high affinity for the various human alpha 1a adrenergic receptors. Furthermore, functional biological tests are used to confirm the effects of identified compounds as alpha 1a adrenergic receptor antagonists.

The present invention also has the objective of providing suitable topical, oral, systemic and parenteral pharmaceutical formulations for use in the novel methods of treatment of the present invention. The compositions containing compounds of this invention as the active ingredient for use in the specific antagonism of human alpha 1a adrenergic receptors can be administered in a wide variety of therapeutic dosage forms in conventional vehicles for systemic administration. For example, the compounds can be administered in such oral dosage forms as tablets, capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, solutions, suspensions, syrups and emulsions, or by injection. Likewise, they may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed as an alpha 1a antagonistic agent.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be

administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

5 The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound thereof employed. A physician or veterinarian of ordinary skill can readily determine and prescribe the effective
10 amount of the drug required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentration of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug.

15 In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups
20 and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants,
25 disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms
30 include, without limitation, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The liquid forms in suitably flavored suspending or dispersing agents
35 such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-

cellulose and the like. Other dispersing agents which may be employed include glycerin and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

5 The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

10 Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinyl-pyrrolidone, pyran copolymer, polyhydroxypropylmethacryl-
15 amidephenol, polyhydroxy-ethylaspartamidephenol, or polyethy-
 leneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyeppilon caprolactone, polyhydroxy butyric acid, polyorthoesters,
20 polyacetals, polydihydro-pyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

 Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever specific blockade of the human alpha 1a adrenergic receptor is required.

25 The daily dosage of the products may be varied over a wide range; e.g., from about 0.01 to about 1000 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0 and 100 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated.
30 A medicament typically contains from about 0.01 mg to about 500 mg of the active ingredient, preferably, from about 1 mg to about 100 mg of active ingredient. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.0002 mg/kg to about 20 mg/kg of body weight per day. Preferably, the range is from about 0.001 to about 10 mg/kg of body weight per day, and especially from
35 about 0.001 mg/kg to about

7 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 4 times per day.

Compounds of this patent disclosure may be used alone at appropriate dosages defined by routine testing in order to obtain optimal antagonism of the human
5 alpha 1a adrenergic receptor while minimizing any potential toxicity. In addition, co-administration or sequential administration of other agents which alleviate the effects of BPH is desirable. Thus, in one embodiment, this invention is administration of compounds of this invention and a human testosterone 5-a reductase inhibitor. Included with this embodiment are inhibitors of 5-alpha reductase isoenzyme 2.
10 Many such compounds are now well known in the art and include such compounds as PROSCAR®, (also known as finasteride, a 4-Aza-steroid; see US 4377584 and 4760071, for example). In addition to PROSCAR®, which is principally active in prostatic tissue due to its selectivity for human 5-a reductase isozyme 2, combinations of compounds which are specifically active in inhibiting testosterone 5-alpha
15 reductase isozyme 1 and compounds which act as dual inhibitors of both isozymes 1 and 2, are useful in combination with compounds of this invention. Compounds that are active as 5a-reductase inhibitors have been described in WO93/23420, EP 0572166;
WO 93/23050; WO93/23038, ; WO93/23048; WO93/23041; WO93/23040;
20 WO93/23039; WO93/23376; WO93/23419, EP 0572165; WO93/23051.

The dosages of the alpha 1a adrenergic receptor and testosterone 5-alpha reductase inhibitors are adjusted when combined to achieve desired effects. As those skilled in the art will appreciate, dosages of the 5-alpha reductase inhibitor and the alpha 1a adrenergic receptor antagonist may be independently
25 optimized and combined to achieve a synergistic result wherein the pathology is reduced more than it would be if either agent were used alone. In accordance with the method of the present invention, the individual components of the combination can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. The instant
30 invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly.

Thus, in one embodiment of the present invention, a method of treating BPH is provided which comprises administering to a subject in need of
35 treatment any of the compounds of the present invention in combination with

finasteride effective to treat BPH. The dosage of finasteride administered to the subject is from about 0.01 mg per subject per day to about 50 mg per subject per day in combination with an alpha 1a antagonist. In one aspect, the dosage of finasteride in the combination is from about 0.2 mg per subject per day to about 10 mg per subject per day, and, in another aspect, from about 1 to about 7 mg per subject to day (e.g., about 5 mg per subject per day).

For the treatment of benign prostatic hyperplasia, compounds of this invention exhibiting alpha 1a adrenergic receptor blockade can be combined with a therapeutically effective amount of a 5a-reductase 2 inhibitor, such as finasteride, in addition to a 5a-reductase 1 inhibitor, such as 4,7B-dimethyl-4-aza-5a-cholestan-3-one, in a single oral, systemic, or parenteral pharmaceutical dosage formulation. Alternatively, a combined therapy can be employed wherein the alpha 1a adrenergic receptor antagonist and the 5a-reductase 1 or 2 inhibitor are administered in separate oral, systemic, or parenteral dosage formulations. See, e.g., US 4377584 and US 4760071 which describe dosages and formulations for 5a-reductase inhibitors.

Abbreviations used in the instant specification, particularly the Schemes and Examples, are as follows:

- DIPEA = diisopropyl ethylamine
- DMF = N,N-dimethylformamide
- DMSO = dimethylsulfoxide
- EDTA = ethylenediamine tetraacetic acid
- Et = ethyl
- Et₂O = diethyl ether
- EtOAc = ethyl acetate
- FAB MS = fast atom bombardment mass spectroscopy
- LDA = lithium diisopropyl amide
- LHMDS = lithium bis(trimethylsilyl)amide
- Me = methyl
- MeOH = methanol
- NMR = nuclear magnetic resonance
- PCTLC = preparative centrifugal thin layer chromatography
- PMBCl = p-methoxybenzyl chloride
- (p-NO₂Ph)COCl = p-nitrophenylchloroformate

TFA = trifluoroacetic acid

THF = tetrahydrofuran


The compounds of the present invention can be readily prepared according to the following reaction schemes and examples, or modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in this art, but are not mentioned in greater detail. Furthermore, other methods for preparing compounds of the invention will be readily apparent to the person of ordinary skill in the art in light of the following reaction schemes and examples. Unless otherwise indicated, all variables are as defined above.

Many of the compounds of the present invention can be prepared via Scheme 1 shown below. In Scheme 1, the ortho-aminoaryl- or o-aminoheteroaryl-nitrile **1** is treated with an arylmagnesium bromide or an aryllithium, followed by diethylcarbonate to afford aryl-substituted polyazanaphthalenone **2** (e.g., 4-aryl-quinazolin-2-one or 4-aryl-pteridin-2-one). Compound **2** is then reacted either with LHMDs and 4-methoxybenzyl chloride or with LHMDs and R¹D (wherein R¹ = C₁-C₈ alkyl or fluorinated C₁-C₈ alkyl, and D is chloro, bromo, iodo, mesylate, tosylate, nosylate, or triflate) to form nitrogen-alkylated analog **3**. Compound **3** is reacted either with R¹Cu (other than R¹ = H) or with NaBH₄ (to give R¹ = H) to form dihydro derivative **4**. Deprotonation of **4** with a strong base (for example, LDA) and addition to a THF solution of p-nitrophenylchloroformate produces stable, isolable "activated" dihydro polyazanaphthalenone **5**, which is coupled with amines of formula Q-H to give coupled product **6**, which for R* = R² (= R² other than H) is a compound of the invention **7a**. For R* = PMB, coupled product **6** is deprotected to afford **7b** (i.e., R² = H).


Methods for preparing the amines of formula Q-H are described in US 5661163, WO 98/57632, WO 98/57638, WO 98/57639, WO 98/57640, WO 98/57641 and WO 98/57642.

Scheme 2 illustrates the preparative procedures employed in Examples 1-6 below.


SCHEME 1




SCHEME 1 (continued)




SCHEME 2



SCHEME 2 (continued)




SCHEME 2 (continued)



The following Examples further describe and illustrate the invention and its practice and are not to be construed as limiting the scope or spirit of the invention.

EXAMPLE 1

N-(3-Amino)propyl-4-(2-cyanophenyl)-4-cyanopiperidine dihydrochloride



Step A: Bis(2-chloroethyl)-*N*-(1,1-dimethylethoxy)carbonyl amine

A solution of *N*-(2,2'-bischloro)diethyl amine (23.0 g, 0.130 mol) and di-*tert*-butyl dicarbonate (28.8 g, 0.130 mol) in CH_2Cl_2 (150 mL) was treated with *N,N*-diisopropylethylamine (22.52 mL, 0.720 mol) at room temperature (1.5 h). The solvent was removed *in vacuo* and the residue was triturated with ether (300 mL). The ether solution was collected and concentrated *in vacuo* affording *N*-(2,2'-bischloro)-diethyl-*N*-(1,1-dimethylethoxy)carbonyl amine as a clear oil.

^1H NMR (CDCl_3): δ 3.65 (m, 8H), 1.52 (s, 9H)

FAB MS (3:1 mixture of dithiothreitol and dithioerythritol in MeOH) m/e 242 g/mole ($\text{M}^+ + \text{H}$, $\text{C}_{25}\text{H}_{29}\text{N}_2\text{O}_5\text{SCl} = 242.2$ g/mole.)

Step B: 4-Cyano-*N*-(1,1-dimethylethoxy)carbonyl-4-(2-cyanophenyl) piperidine

A solution of bis(2-chloroethyl)-*N*-(1,1-dimethylethoxy)carbonyl amine (19 g, 78 mmol) and homophthalonitrile (8.6 g, 60 mmol) and cesium carbonate (79 g, 241 mmol) was stirred at 60 °C for 12 hours. The solution cooled to room temperature and diluted with 500 ml EtOAc and washed with saturated aqueous NaHSO₄ and saturated aqueous NaCl. The solution was dried over Na₂SO₄, concentrated *in vacuo*, and purified by PCTLC (30% EtOAc in Hexane) to afforded 4-cyano-*N*-(1,1-dimethylethoxy)carbonyl-4-(2-cyanophenyl) piperidine as a yellow/orange oil.

¹H NMR (CDCl₃): δ 7.81-7.79 (d, 2H), 7.67-7.65 (m, 2H), 7.51-7.47 (m, 1H), 4.31 (br s, 2H), 3.26 (br s, 2H), 2.33-2.20 (m, 4H), 1.48 (s, 9H).

Step C: 4-Cyano-4-(2-cyanophenyl)piperidine hydrochloride

15

A solution of EtOAc saturated with HCl (200 ml) was added to 4-cyano-*N*-(1,1-dimethylethoxy)carbonyl-4-(2-methylphenyl)piperidine (1.5 g, 5.7 mmol). The resulting mixture was allowed to react for 1 hour at room temperature. The EtOAc was removed *in vacuo* affording 1.5 g of 4-cyano-4-(2-cyanophenyl)piperidine hydrochloride as a yellow solid.

20

Step D: 4-(2-Cyanophenyl)-4-cyano-*N*-(3-[*N*-(1,1-dimethyl ethoxycarbonyl)] amino)propylpiperidine

25

A solution of 4-(2-cyanophenyl)-4-cyanopiperidine hydrochloride (0.5 g, 2.01 mmol) and *N,N*-diisopropylethylamine (0.421 ml, 2.42 mmol) in DMF (5 mL) at 0°C was added sodium hydride (0.0968 g, 2.42 mmol). The slurry stirred for 1 hour at 0°C and was then added *N*-(3-bromopropyl)-*N*-(1,1-dimethylethoxy)carbonyl amine (0.480 g, 2.01 mmol) in DMF (3 mL). The solution stirred for 12 hours at room temperature. The reaction was quenched by the addition of water. The product was extracted with EtOAc (2 x 50 ml) and washed with H₂O (2 x 25 ml). The solvent was dried over Na₂SO₄, concentrated *in vacuo*, and purified by PCTLC (5% MeOH in CHCl₃). 4-(2-Cyanophenyl)-4-cyano-*N*-(3-[*N*-(1,1-dimethyl-ethoxy)carbonyl] amino)propyl piperidine was obtained as an oil.

30

¹H NMR (CDCl₃): δ 7.80-7.78 (dd, 1H), 7.67-7.59 (m, 2H), 7.49-7.45 (m, 1H), 5.14 (br s, 1H), 3.21-3.20 (d, 2H), 3.09-3.06 (d, 2H), 2.58-2.51 (m, 3H), 2.43-2.40 (d, 2H), 2.30-2.25 (m, 2H), 1.73-1.66 (m, 2H), 1.56 (s, 1H), 1.43 (s, 9H).


5 Step E: *N*-(3-Amino)propyl-4-(2-cyanophenyl)-4-cyanopiperidine dihydrochloride

The title compound was prepared using the same procedure described above in Step C for the preparation of 4-cyano-4-(2-cyanophenyl)piperidine hydrochloride.

10

EXAMPLE 2

3-[4-(4-Fluorophenyl)piperidin-1-yl]propylamine



15 Step A. 4-(4-Fluorophenyl)piperidine hydrochloride

To a solution of 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine hydrochloride (10 g) in methanol (200 mL) was added 10% palladium on charcoal (0.5 g) and the mixture was hydrogenated at 50 psi for 3 h. The catalyst was removed
20 by filtration and solvent was evaporated to leave the product as a white powder, which was used in the next step without any purification. C

M.P. 181-182 °C.

¹H NMR (CDCl₃): δ 1.95-2.03 (br d, 2H), 2.14-2.29 (m, 2H), 2.70-2.80 (m, 1H), 2.91-3.07 (br q, 2H), 3.60-3.64 (br d, 2H), 6.96-7.03 (m, 2H), 7.19-7.22
25 (m, 2H), 9.60 (br s, 1H), 9.71 (br s, 1H).

Step B. 3-[4-(4-Fluorophenyl)piperidin-1-yl]propylphthalimide

A mixture of 4-(4-fluorophenyl)piperidine hydrochloride (5.08 g, 23.2 mmol), 3-bromopropylphthalimide (6.22 g, 23.2 mmol), and potassium carbonate (15 g) in DMF (100 mL) was stirred and heated at 95-100 °C for 12 h. About 80% of the solvent was evaporated at reduced pressure, the residue was diluted with ethyl acetate

(200 mL) and washed with brine (3 X 100 mL) and dried (Na₂SO₄). Solvent was evaporated and the residue was purified by column chromatography on silica gel using 1/1 hexane-ethyl acetate to 100% ethyl acetate as eluent. This product was crystallized from isopropanol to give a white crystalline solid; m.p. 80-81 °C. This material was used in the next step. Concentration of the mother liquor and cooling gave the second crop.

¹H NMR (CDCl₃): δ 1.43-1.52 (m, 2H), 1.67-1.75 (m, 2H), 1.80-1.96 (m, 4H), 2.33-2.46 (m, 3H), 2.94-2.99 (br d, 2H), 3.78 (t, *J* = 7 Hz, 2H), 6.90-7.04 (m, 4H), 7.70-7.74 (m, 2H), 7.84-7.87 (m, 2H).


Step C. 3-[4-(4-Fluorophenyl)piperidin-1-yl]propylamine

To a solution of 3-[4-(4-fluorophenyl)piperidin-1-yl]propylphthalimide (4.5 g, 12.3 mmol) in methanol (200 mL) was added 4 ml of hydrazine and the mixture was refluxed for 8 h. It was cooled, and the white solid was filtered and washed with methanol (20 mL). Solvent was evaporated, and the residue was dried under vacuum for 4 h. Chloroform (50 mL) was added to this material, it was stirred for 1 h and filtered. The white solid was washed with more chloroform (20 mL), and the solvent was evaporated from the combined filtrates to leave the crude product as an oil. It was purified by column chromatography on silica gel using dichloromethane/methanol/2M ammonia in methanol (10/3/1) as the eluent.

¹H NMR (CDCl₃): δ 1.60-1.83 (m, 6H), 1.96-2.07 (m, 4H), 2.40-2.55 (m, 3H), 2.70-2.85 (br t, 2H), 3.03-3.07 (br d, 2H), 6.93-7.00 (m, 2H), 7.14-7.20 (m, 2H).

EXAMPLE 3


4-(3,4-Difluorophenyl)-quinazolin-2-one (8)



- 5 An oven dried round-bottom flask was charged with magnesium turnings (3.55 g, 146 mmol) in THF (100 mL) under argon. 1-Bromo-3,4-difluorobenzene (4.0 mL, 35.3 mmol) was added to the suspension in one portion. The solution was heated to 40 °C. Once the temperature had stabilized at 50 °C, the remainder of the bromide (13.52 mL, 155 mmol total) was added dropwise. After
- 10 complete addition, the solution was stirred for 1 hour at 65 °C. A solution of anthranilonitrile (4.95 g, 41.9 mmol) in THF (25 mL) was added dropwise while keeping the internal temperature constant. After stirring or two hours at 65 °C, diethyl carbonate (16.3 mL, 135 mmol) was added dropwise to the brownish-red solution. This solution was stirred for an additional 30 minutes at 65 °C and then
- 15 cooled to room temperature. The solution was poured into a mixture of 250 mL of 1M citric acid and ice. The mixture was extracted three times with CHCl₃ (50 mL) and the combined extracts were washed with a 10% Na₂CO₃ solution. The organics were dried over Na₂SO₄, concentrated, and triturated with Et₂O to afford the title compound.
- 20 ¹H NMR (DMSO-d₆, 400 MHz) 7.81-7.72 (m, 2H), 7.69-7.62 (m, 2H), 7.57-7.53 (m, 1H), 7.38-7.36 (d, 1H), 7.23-7.19 (t, 1H).

EXAMPLE 4

4-(3,4-Difluorophenyl)-1-(4-methoxybenzyl)-quinazolin-2-one (9)



5 To a 0 °C solution of 8 (9.73 g, 37.7 mmol) in DMF (100 mL) under argon was added lithium bis(trimethylsilyl) amide (43.9 mL, 1.0 M) dropwise. 4-Methoxybenzyl chloride (8.52 mL, 62.8 mmol) was then added in one portion. The solution was stirred at 55 °C for sixteen hours. The solvent was removed *in vacuo* and the residue was partitioned between CHCl₃ and cold 1.0 M citric acid. The


10 aqueous layer was extracted with CHCl₃, and the combined organic extracts were washed with 10% Na₂CO₃ and dried. The crude product was not purified. A light yellow solid was obtained.

¹H NMR (DMSO-d₆, 400 MHz) 7.86-7.64 (m, 4H), 7.59-7.57 (d, 2H), 7.30-7.27 (m, 3H), 6.90-6.88 (d, 2H), 5.46 (s, 2H), 3.71 (s, 3H).

15

EXAMPLE 5

4-(3,4-Difluorophenyl)-1-(4-methoxybenzyl)-3,4-dihydro-quinazolin-2-one (10)




To a 0 °C solution of 9 (9.33 g, 24.6 mmol) in dioxane (250 mL) was added NaBH₄ (1.02 g, 27.1 mmol) in one portion. The solution stirred for two hours at room temperature. The solvent was removed *in vacuo*, and the residue was partitioned between CHCl₃ and 1 N HCl. The aqueous layer was extracted with CHCl₃, and the combined organic extracts were dried, concentrated, and purified by PCTLC (1:1 EtOAc:Hex). A white solid was obtained.

¹H NMR (CDCl₃, 400 MHz) 7.30-7.04 (m, 6H), 6.93-6.83 (m, 6H), 5.58 (s, 1H), 5.18-5.04 (q, 2H), 3.77 (s, 3H).

EXAMPLE 6

4-(3,4-Difluorophenyl)-1-(4-methoxybenzyl)-3-(4-nitrophenoxycarbonyl)-3,4-dihydro-quinazolin-2-one (11)




The title compound was prepared by treating **10** (0.292 g, 0.767 mmol) with lithium diisopropylamide (2.0M THF solution, 1.1 equivalents) in THF at -78 °C for 20 minutes followed by the rapid addition of 4-nitrophenyl chloroformate (1.5 equivalents) in THF. The ¹H NMR was consistent with the assigned structure.

5

EXAMPLE 7

4-(3,4-Difluorophenyl)-1-(4-methoxybenzyl)-3-((4-cyano-4-(2-cyanophenyl)piperidin-1-yl)propylcarbamoyl)-3,4-dihydro-quinazolin-2-one (**12**)



10

To a solution of **11** (0.060 g, 0.109 mmol) in 10 ml of CHCl₃ was added *N*-(3-amino)propyl-4-(2-cyanophenyl)-4-cyanopiperidine dihydrochloride (Example 1) and diisopropyl ethylamine (0.057 mL, 0.329 mmol). The resulting solution was stirred for 2 hours and the crude material purified by PCTLC (5% MeOH in CHCl₃/2%NH₄OH) affording the title compound.

15

¹H NMR (CDCl₃, 400 MHz) 9.10-9.07 (t, 1H), 7.78-7.76 (d, 1H), 7.54-7.48 (m, 3H), 7.13-7.09 (t, 1H), 7.02-6.93 (q, 1H), 6.91-6.83 (m, 5H), 6.75-6.73 (d, 2H), 5.05-5.01 (d, 1H), 4.88-4.84 (d, 2H), 3.77 (s, 3H) 3.49-3.45 (m, 3H), 3.12-3.03 (m, 2H), 2.57-2.51 (m, 4H), 2.38-2.35 (d, 3H), 2.26-2.23 (m, 1H), 1.82-1.77 (m, 2H), 1.53 (s, 2H).

20


MS (FAB) 675.4 (M+1)

Analysis calculated for C₃₉H₃₆N₆O₃F₂: C:69.42, H:5.37, N:12.45,

Found C:69.43, H:5.36, N:12.14.

EXAMPLE 8

4-(3,4-Difluorophenyl)-3-((4-cyano-4-(2-cyanophenyl)piperidin-1-yl)propylcarbamoyl)-3,4-dihydro-quinazolin-2-one (13)



5 A solution of 12 (0.050 g, 0.074 mmol) in TFA (1.5 mL) was stirred for six hours at 65 °C. The solvent was removed *in vacuo* and azeotroped twice with toluene. The product was then purified by PCTLC (5% MeOH in CHCl₃/2%NH₄OH). A white solid was obtained.

¹H NMR (CDCl₃, 400 MHz) 9.06-9.04 (t, 1H), 7.79-7.77 (d, 1H),
10 7.66-7.58 (m, 2H), 7.48-7.44 (t, 1H), 7.40 (s, 1H), 7.30-7.23 (m, 2H), 7.11-6.98 (m, 4H), 6.89 (s, 1H), 6.85-6.83 (d, 1H), 3.46-3.37 (m, 2H) 3.07-3.04 (m, 2H), 2.58-2.51 (m, 4H), 2.40-2.27 (m, 4H), 1.81-1.76 (m, 2H).


MS (FAB) 555.2 (M+1)

Analysis calculated for C₃₁H₂₈N₆O₂F₂·50 EtOAc: C:66.20, H:5.39,

15 N:14.04, Found C:65.87, H:5.15, N:14.07.

EXAMPLE 9

4-(3,4-Difluorophenyl)-1-(4-methoxybenzyl)-3-(4-(4-fluorophenyl)piperidin-1-yl)propylcarbamoyl)-3,4-dihydro-quinazolin-2-one (14)



- 5 The title compound was prepared using the same procedure described for the preparation of 12, except that 3-[4-(4-fluorophenyl)piperidin-1-yl]propylamine (Example 2) was used in place of *N*-(3-amino)propyl-4-(2-cyanophenyl)-4-cyanopiperidine dihydrochloride.

- 10 ¹H NMR (CDCl₃, 400 MHz) 9.03-9.00 (t, 1H), 7.30-7.24 (q, 2H), 7.17-7.09 (m, 3H), 7.04-6.91 (m, 7H), 6.88-6.82 (m, 2H), 6.76-6.74 (d, 2H), 5.17-5.13 (d, 1H), 4.95-4.91 (d, 1H), 3.45-3.41 (q, 2H) 3.05-3.03 (m, 2H), 2.46-2.42 (t, 3H), 2.03-1.97 (m, 2H), 1.82-1.76 (m, 6H).


MS (FAB) 643.3 (M+1)

Analysis calculated for C₃₇H₃₇N₄O₃F₂.60 HCl + 1.05 CHCl₃ :

- 15 C:57.85, H:4.93, N:7.09, Found C:57.86, H:5.31, N:6.71.

EXAMPLE 10

4-(3,4-Difluorophenyl)-3-(4-(4-fluorophenyl)piperidin-1-yl)propylcarbamoyl)-3,4-dihydro-quinazolin-2-one (15)



5 The title compound was prepared using the same procedure described for the preparation of 13.


¹H NMR (CDCl₃, 400 MHz) 9.04-9.01 (t, 1H), 7.55 (s, 1H), 7.31-7.24 (m, 2H), 7.17-7.02 (m, 5H), 7.00-6.94 (m, 3H), 6.90-6.83 (m, 2H), 3.44-3.37 (m, 2H) 3.05-3.02 (d, 2H), 2.47-2.42 (m, 3H), 2.07-2.00 (m, 2H), 1.82-1.73 (m, 6H).

10 MS (FAB) 523.3 (M+1)

Analysis calculated for C₂₉H₂₉N₄O₂F₃ 1.45 HCl + .35 EtOAc:
C:60.22, H:5.53, N:9.24, Found C:60.21, H:5.64, N:9.32.

EXAMPLE 11

15 *trans*-4-(3,4-Difluorophenyl)-3-(1-(4-(2-cyanophenyl)cyclohexyl)-3-hydroxy-azetidin-3-yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2-one (16)



The title compound was prepared using a procedure analogous to that described for the preparation of 13, except that *trans*-1-(4-(2-

cyanophenyl)cyclohexyl)-3-hydroxy-azetidin-3-yl)methylamine was used in place of *N*-(3-amino)propyl-4-(2-cyanophenyl)-4-cyanopiperidine dihydrochloride. 1-(4-(2-Cyanophenyl)cyclohexyl)-3-hydroxy-azetidin-3-yl)methylamine was prepared in accordance with procedures described in WO 98/57641.


¹H NMR (CDCl₃, 400 MHz) 9.58-9.55 (t, 1H), 8.83 (br s, 1 H), 7.63-7.61 (d, 1H), 7.57-7.53 (t, 1H), 7.37-7.35 (d, 1H), 7.31-7.26 (m, 3H), 7.23-7.21 (d, 1H), 7.09-7.01 (m, 3H), 6.98-6.91 (m, 2H), 6.84 (s, 1H) 3.62-3.61 (d, 2H) 3.41-3.37 (m, 2H), 3.29-3.25 (m, 2H), 3.06-3.00 (t, 1H), 2.18 (s, 1H), 1.98-1.88 (m, 4H), 1.51-1.43 (m, 2H), 1.36-1.33 (m, 2H).

MS (FAB) 572.1 (M+1)

Analysis calculated for C₃₂H₃₁N₅O₃F₂.15 CH₂Cl₂ + .05 Et₂O
C:66.07, H:5.45, N:11.91, Found C:65.95, H:5.37, N:11.87.

EXAMPLE 12

trans-4-(3,4-Difluorophenyl)-3-(1-(4-(2-cyanophenyl)cyclohexyl)azetidin-3-yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2-one (17)



The title compound was prepared using a procedure analogous to that described for the preparation of 13, except that *trans*-1-(4-(2-cyanophenyl)cyclohexyl)azetidin-3-ylmethylamine was used in place of *N*-(3-amino)propyl-4-(2-cyanophenyl)-4-cyanopiperidine dihydrochloride. 1-(4-(2-Cyanophenyl)cyclohexyl) azetidin-3-ylmethylamine was prepared in accordance with procedures described in WO 98/57641.

¹H NMR (CDCl₃, 400 MHz) 9.15-9.12 (t, 1H), 7.90 (br s, 1 H), 7.60-7.58 (d, 1H), 7.54-7.50 (t, 1H), 7.33-7.23 (m, 4H), 7.12-6.96 (m, 4H), 6.89-6.85 (t, 2H), 3.53-3.48(m, 2H) 3.41-3.38 (t, 2H), 2.98-2.87 (m, 3H), 2.72-2.68 (m, 1H), 2.07-2.01 (m, 1H), 1.92-1.90 (m, 4H), 1.54-1.45 (m, 2H), 1.22-1.17 (m, 2H).

MS (FAB) 556.1 (M+1)


Analysis calculated for $C_{32}H_{31}N_5O_2F_{2.35}CH_2Cl_2 + .20 Et_2O$

C:66.34, H:5.66, N:11.67, Found C:66.47, H:5.61, N:11.66.

5

EXAMPLE 13

trans-4-(3,4-Difluorophenyl)-3-(1-(4-(2-pyridyl)cyclohexyl)azetidin-3-yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2-one (18)



The title compound was prepared using a procedure analogous to that described for the preparation of 13, except that *trans*-1-(4-(2-pyridyl)cyclohexyl)azetidin-3-ylmethyl amine was used in place of *N*-(3-amino)propyl-4-(2-cyanophenyl)-4-cyanopiperidine dihydrochloride. 1-(4-(2-Pyridyl)cyclohexyl)azetidin-3-ylmethyl amine was prepared in accordance with procedures described in WO 98/57641.

1H NMR (CDCl₃, 400 MHz) 9.13-9.1 (t, 1H), 8.52-8.50 (d, 1H), 8.00 (br s, 1H), 7.60-7.56 (m, 1H), 7.30-7.22 (m, 2H), 7.13-6.98 (m, 6H), 6.87-6.84 (t, 2H), 3.54-3.48 (m, 2H), 3.42-3.38 (t, 2H), 2.98-2.94 (m, 3H), 2.72-2.61 (m, 2H), 2.04-1.88 (m, 5H), 1.56-1.53 (q, 2H), 1.14-1.10 (m, 2H).

MS (FAB) 532.1 (M+1)

Analysis calculated for $C_{30}H_{31}N_5O_2F_{2.45}CH_2Cl_2 + .25 Et_2O$
C:64.20, H:5.89, N:11.90, Found C:64.18, H:5.71, N:11.91.

EXAMPLE 14

As a specific embodiment of an oral composition, 100 mg of the compound of Example 10 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

EXAMPLE 15

Screening assay: Alpha 1a Adrenergic Receptor Binding

Membranes prepared from the stably transfected human alpha 1a cell line (ATCC CRL 11140) were used to identify compounds that bind to the human alpha 1a adrenergic receptor. These competition binding reactions (total volume = 200 μ l) contained 50 mM Tris-HCl pH. 7.4, 5 mM EDTA, 150 mM NaCl, 100 pM [125 I]-HEAT, membranes prepared from the alpha 1a cell line and increasing amounts of unlabeled ligand. Reactions were incubated at room temperature for one hour with shaking. Reactions were filtered onto Whatman GF/C glass fiber filters with a Inotec 96 well cell harvester. Filters were washed three times with ice cold buffer and bound radioactivity was determined (K_i).

EXAMPLE 16

Selective Binding assays

Membranes prepared from stably transfected human alpha 1d and alpha 1b cell lines (ATCC CRL 11138 and CRL 11139, respectively) were used to identify compounds that selectively bind to the human alpha 1a adrenergic receptor. These competition binding reactions (total volume = 200 μ l) contained 50 mM Tris-HCl pH. 7.4, 5 mM EDTA, 150 mM NaCl, 100 pM [125 I]-HEAT, membranes prepared from cell lines transfected with the respective alpha 1 subtype expression plasmid and increasing amounts of unlabeled ligand. Reactions were incubated at room temperature for one hour with shaking. Reactions were filtered onto Whatman GF/C glass fiber filters with a Inotec 96 well cell harvester. Filters were washed three times with ice cold buffer and bound radioactivity was determined (K_i).

The compounds of the present invention prepared in the foregoing examples were found to have alpha 1a K_i values of less than about 20 nM as determined via the screening assay described in Example 15. All of the compounds were further found to be at least about 15-fold more selective in binding to alpha 1a receptors versus binding to the alpha 1b and alpha 1d receptors, as determined via the selective binding assay described in the preceding paragraph.

EXAMPLE 17

Counterscreen: Histamine-1 Selectivity

The binding affinity (K_i in nM) of the compounds of the present invention for histamine H1 receptors can be determined via the binding assay described

in Chang et al., *J. Neurochem.* (1979), **32**: 1653, or as described in US 5403847, or suitable modifications thereof known to those skilled in the art. The assay can be used to eliminate agents which specifically affect binding to hH1 receptors.

5 EXAMPLE 18

EXEMPLARY COUNTERSCREENS

1. Assay Title: Dopamine D2, D3, D4 in vitro screen

10 Objective of the Assay:

The objective of this assay is to eliminate agents which specifically affect binding of [^3H] spiperone to cells expressing human dopamine receptors D2, D3 or D4.

15 Method:

Modified from VanTol et al., *Nature* (1991), 350: 610-613.

Frozen pellets containing specific dopamine receptor subtypes stably expressed in clonal cell lines are lysed in 2 ml lysing buffer (10mM Tris-HCl/5mM Mg, pH 7.4). Pellets obtained after centrifuging these membranes (15' at 24,450 rpm) are resuspended in 50mM Tris-HCl pH 7.4 containing EDTA, MgCl₂, KCl, NaCl, CaCl₂ and ascorbate to give a 1 Mg/mL suspension. The assay is initiated by adding 50-75 µg membranes in a total volume of 500 µl containing 0.2 nM [³H]-spiperone. Non-specific binding is defined using 10 µM apomorphine. The assay is terminated after a 2 hour incubation at room temperature by rapid filtration over GF/B filters presoaked in 0.3% PEI, using 50mM Tris-HCl pH 7.4.

2. Assay Title: Serotonin 5HT1a

Objective of the Assay

30 The objective of this assay is to eliminate agents which specifically
affect binding to cloned human 5HT1a receptor

Method:

Modified from Schelegel and Peroutka, *Biochemical Pharmacology* 35 (1986), 35: 1943-1949.

Mammalian cells expressing cloned human 5HT_{1a} receptors are lysed in ice-cold 5 mM Tris-HCl, 2 mM EDTA (pH 7.4) and homogenized with a polytron homogenizer. The homogenate is centrifuged at 1000Xg for 30', and then the supernatant is centrifuged again at 38,000Xg for 30'. The binding assay contains 0.25 nM [³H]8-OH-DPAT (8-hydroxy-2-dipropylamino-1,2,3,4-tetrahydronaphthalene) in 50 mM Tris-HCl, 4 mM CaCl₂ and 1mg/ml ascorbate. Non-specific binding is defined using 10 μM propranolol. The assay is terminated after a 1 hour incubation at room temperature by rapid filtration over GF/C filters

EXAMPLE 19

EXEMPLARY FUNCTIONAL ASSAYS

In order to confirm the specificity of compounds for the human alpha 1a adrenergic receptor and to define the biological activity of the compounds, the following functional tests may be performed:

1. In vitro Rat, Dog and Human Prostate and Dog Urethra

Taconic Farms Sprague-Dawley male rats, weighing 250-400 grams are sacrificed by cervical dislocation under anesthesia (methohexital; 50 mg/kg, i.p.). An incision is made into the lower abdomen to remove the ventral lobes of the prostate. Each prostate removed from a mongrel dog is cut into 6-8 pieces longitudinally along the urethra opening and stored in ice-cold oxygenated Krebs solution overnight before use if necessary. Dog urethra proximal to prostate is cut into approximately 5 mm rings, the rings are then cut open for contractile measurement of circular muscles. Human prostate chips from transurethral surgery of benign prostate hyperplasia are also stored overnight in ice-cold Krebs solution if needed.

The tissue is placed in a Petri dish containing oxygenated Krebs solution [NaCl, 118 mM; KCl, 4.7 mM; CaCl₂, 2.5 mM; KH₂PO₄, 1.2 mM; MgSO₄, 1.2 mM; NaHCO₃, 2.0 mM; dextrose, 11 mM] warmed to 37°C. Excess lipid material and connective tissue are carefully removed. Tissue segments are attached to glass tissue holders with 4-0 surgical silk and placed in a 5 ml jacketed tissue bath containing Krebs buffer at 37 °C, bubbled with 5% CO₂/95% O₂. The tissues are connected to a Satham-Gould force transducer; 1 gram (rat, human) or 1.5 gram

(dog) of tension is applied and the tissues are allowed to equilibrate for one hour. Contractions are recorded on a Hewlett-Packard 7700 series strip chart recorder.

After a single priming dose of 3 μ M (for rat), 10 μ M (for dog) and 20 μ M (for human) of phenylephrine, a cumulative concentration response curve to an agonist is generated; the tissues are washed every 10 minutes for one hour. Vehicle or antagonist is added to the bath and allowed to incubate for one hour, then another cumulative concentration response curve to the agonist is generated.

EC₅₀ values are calculated for each group using GraphPad Inplot software. pA₂ (-log K_b) values were obtained from Schild plot when three or more concentrations were tested. When less than three concentrations of antagonist are tested, K_b values are calculated according

to the following formula $K_b = \frac{[B]}{x-1}$,

$$x-1$$

where x is the ratio of EC₅₀ of agonist in the presence and absence of antagonist and [B] is the antagonist concentration.

2. Measurement of Intra-Urethral Pressure in Anesthetized Dogs

PURPOSE: Benign prostatic hyperplasia causes a decreased urine flow rate that may be produced by both passive physical obstruction of the prostatic urethra from increased prostate mass as well as active obstruction due to prostatic contraction. Alpha adrenergic receptor antagonists such as prazosin and terazosin prevent active prostatic contraction, thus improve urine flow rate and provide symptomatic relief in man. However, these are non-selective alpha 1 receptor antagonists which also have pronounced vascular effects. Because we have identified the alpha 1a receptor subtype as the predominant subtype in the human prostate, it is now possible to specifically target this receptor to inhibit prostatic contraction without concomitant changes in the vasculature. The following model is used to measure adrenergically mediated changes in intra-urethral pressure and arterial pressure in anesthetized dogs in order to evaluate the efficacy and potency of selective alpha adrenergic receptor antagonists. The goals are to: 1) identify the alpha-1 receptor subtypes responsible for prostatic/urethral contraction and vascular responses, and 2) use this model to evaluate novel selective alpha adrenergic antagonists. Novel and standard alpha adrenergic antagonists may be evaluated in this manner.

METHODS: Male mongrel dogs (7-12 kg) are used in this study. The dogs are anesthetized with pentobarbital sodium (35 mg/kg, i.v. plus 4 mg/kg/hr iv infusion). An endotracheal tube is inserted and the animal ventilated with room air using a Harvard instruments positive displacement large animal ventilator. Catheters (PE 240 or 260) are placed in the aorta via the femoral artery and vena cava via the femoral veins (2 catheters, one in each vein) for the measurement of arterial pressure and the administration of drugs, respectively. A supra-pubic incision ~1/2 inch lateral to the penis is made to expose the urethers, bladder and urethra. The urethers are ligated and cannulated so that urine flows freely into beakers. The dome of the bladder is retracted to facilitate dissection of the proximal and distal urethra. Umbilical tape is passed beneath the urethra at the bladder neck and another piece of umbilical tape is placed under the distal urethra approximately 1-2 cm distal to the prostate. The bladder is incised and a Millar micro-tip pressure transducer is advanced into the urethra. The bladder incision is sutured with 2-0 or 3-0 silk (purse-string suture) to hold the transducer. The tip of the transducer is placed in the prostatic urethra and the position of the Millar catheter is verified by gently squeezing the prostate and noting the large change in urethral pressure.


Phenylephrine, an alpha 1 adrenergic agonist, is administered (0.1-100 ug/kg, iv; 0.05 ml/kg volume) in order to construct dose response curves for changes in intra-urethral and arterial pressure. Following administration of increasing doses of an alpha adrenergic antagonist (or vehicle), the effects of phenylephrine on arterial pressure and intra-urethral pressure are re-evaluated. Four or five phenylephrine dose-response curves are generated in each animal (one control, three or four doses of antagonist or vehicle). The relative antagonist potency on phenylephrine induced changes in arterial and intra-urethral pressure are determined by Schild analysis. The family of averaged curves are fit simultaneously (using ALLFIT software package) with a four parameter logistic equation constraining the slope, minimum response, and maximum response to be constant among curves. The dose ratios for the antagonist doses (rightward shift in the dose-response curves from control) are calculated as the ratio of the ED₅₀'s for the respective curves. These dose-ratios are then used to construct a Schild plot and the K_b (expressed as ug/kg, iv) determined. The K_b (dose of antagonist causing a 2-fold rightward shift of the phenylephrine dose-response curve) is used to compare the relative potency of the antagonists on inhibiting phenylephrine responses for intra-urethral and arterial pressure. The

relative selectivity is calculated as the ratio of arterial pressure and intra-urethral pressure K_b 's. Effects of the α_1 antagonists on baseline arterial pressure are also monitored. Comparison of the relative antagonist potency on changes in arterial pressure and intra-urethral pressure provide insight as to whether the α receptor
5 subtype responsible for increasing intra-urethral pressure is also present in the systemic vasculature. According to this method, one is able to confirm the selectivity of α_1 adrenergic receptor antagonists that prevent the increase in intra-urethral pressure to phenylephrine without any activity at the vasculature.

10 While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, the practice of the invention encompasses all of the usual variations, adaptations and/or modifications that come within the scope of the following claims.


WHAT IS CLAIMED IS:

1. A compound of formula:



5

wherein Q is selected from the group consisting of:



A¹, A², A³ and A⁴ are each independently selected from C-X² and N, provided that no more than two of A¹, A², A³ and A⁴ are N, and further provided that when two of A¹, A², A³ and A⁴ are N, each nitrogen atom is bonded to two carbon atoms;

5

each X¹ is independently hydrogen, halogen, cyano, nitro, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl, or (CH₂)₀₋₄OR^a;

each X² is independently hydrogen, halogen, cyano, nitro, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl, or (CH₂)₀₋₄OR^a;

10

R¹ is hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl, phenyl, or substituted phenyl, wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, nitro, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl, fluorinated C₃-C₈ cycloalkyl and (CH₂)₀₋₄OR^a;

15

R² is hydrogen, C₁-C₈ alkyl, or fluorinated C₁-C₈ alkyl;

R³ is hydrogen, C₁-C₈ alkyl, or fluorinated C₁-C₈ alkyl;

20

Y is carbon or nitrogen, provided that when Y is nitrogen, R⁵ is absent;

Z is CH₂, CHOR^b, CHF, CHR^b, C(R^b)₂, CF₂, CHCHF₂, C=CF₂, or C=O;

25

E, G, L and M are each independently selected from hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, (CH₂)₀₋₄OR^c, (CH₂)₀₋₄N(R^c)₂, (CH₂)₀₋₄CN, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂R^c and (CH₂)₀₋₄SO₂N(R^c)₂;

30

J is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, (CH₂)₁₋₄OR^c, (CH₂)₁₋₄N(R^c)₂, (CH₂)₁₋₄CN, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂R^c, or (CH₂)₀₋₄SO₂N(R^c)₂;

R^4 is phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, thienyl, furanyl, substituted pyridyl, substituted thienyl, or substituted furanyl; wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, OR^c, (CH₂)₀₋₃CON(R^c)₂, (CH₂)₀₋₃CO₂R^c, methylenedioxy when the phenyl ring is di-substituted and the substituents are on adjacent carbon atoms, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; and wherein the substituents on the substituted naphthyl, pyridyl, thienyl, or furanyl are independently selected from phenyl, OR^c, halogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl, C₃-C₈ cycloalkyl and fluorinated C₃-C₈ cycloalkyl;

R^5 is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, tetrazole, isooxadiazole, phenyl, or substituted phenyl; wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, (CH₂)₀₋₃CON(R^c)₂, (CH₂)₀₋₃CO₂R^c, methylenedioxy when the phenyl ring is di-substituted and the substituents are on adjacent carbon atoms, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

R^6 and R^7 are each independently selected from hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₁-C₈ alkyl and fluorinated C₃-C₈ cycloalkyl;

R^8 is phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, substituted pyridyl, pyridyl N-oxide (N⁺O⁻), substituted pyridyl N-oxide, pyrazinyl, substituted pyrazinyl, thienyl, substituted thienyl, thiazolyl, substituted thiazolyl, furanyl, substituted furanyl, quinazolinyl, or substituted quinazolinyl; wherein the substituents on the phenyl are independently selected from halogen, cyano, nitro, OR^c, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; and wherein the substituents on the substituted naphthyl, pyridyl, pyrazinyl, thienyl, thiazolyl, furanyl, or quinazolinyl are independently selected from cyano, nitro, N(R^c)₂, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, phenyl, OR^c, halogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl, C₃-C₈ cycloalkyl and fluorinated C₃-C₈ cycloalkyl;

R^9 is hydrogen, C₁-C₈ alkyl, or fluorinated C₁-C₈ alkyl;

R^{10} is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, tetrazole, isooxadiazole, phenyl, substituted phenyl, naphthyl, substituted naphthyl, pyridyl, substituted pyridyl, thienyl, substituted thienyl, furanyl, or substituted furanyl; wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, nitro, OR^c , $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $N(R^c)_2$, NR^cCOR^c , $NR^cCON(R^c)_2$, $NR^cSO_2R^c$, $NR^cSO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl; and wherein the substituents on the substituted naphthyl, pyridyl, thienyl, or furanyl are independently selected from $(CH_2)_{0-4}CO_2R^c$, $(CH_2)_{0-4}CON(R^c)_2$, $(CH_2)_{0-4}SO_2N(R^c)_2$, $(CH_2)_{0-4}SO_2R^c$, phenyl, OR^c , halogen, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl and C₃-C₈ cycloalkyl;

R^{13} , R^{14} , R^{15} and R^{16} are each independently selected from hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, fluorinated C₃-C₈ cycloalkyl, and $(CH_2)_2-4OR^c$;

R^{18} and R^{20} are each independently selected from hydrogen and OR^d ;

R^a is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, or C₃-C₈ cycloalkyl;

R^b is C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or fluorinated C₃-C₈ cycloalkyl;

R^c is hydrogen, C₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or fluorinated C₃-C₈ cycloalkyl;

R^d is hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, or $(CH_2)_{0-4}CF_3$;

m, n, o, and p are each independently integers from 0 to 3;

q₁ is an integer from 0 to 5;

t is an integer from 2 to 5;

u and v are each independently integers from 0 to 3; provided that u and v are not both zero; and further provided that when $u = 0$, Z is selected from CH_2 , CHF , CHR^b , $\text{C(R}^b)_2$, CF_2 , CHCHF_2 and C=CF_2 ; and

5 w is an integer from 0 to 3, provided that when w is 0, R^{20} is hydrogen;

or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1, wherein

10

each X^1 is independently hydrogen, halogen, cyano, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, fluorinated C_1 - C_4 alkyl, fluorinated C_3 - C_6 cycloalkyl, or $(\text{CH}_2)_{0-4}\text{OR}^a$;

each X^2 is independently hydrogen, halogen, cyano, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl,
15 fluorinated C_1 - C_4 alkyl, fluorinated C_3 - C_6 cycloalkyl, or $(\text{CH}_2)_{0-4}\text{OR}^a$;

R^1 is hydrogen, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, fluorinated C_1 - C_4 alkyl, fluorinated
 C_3 - C_6 cycloalkyl, phenyl, or substituted phenyl, wherein the substituents on the
substituted phenyl are independently selected from halogen, cyano, C_1 - C_4 alkyl,
20 fluorinated C_1 - C_4 alkyl and $(\text{CH}_2)_{0-4}\text{OR}^a$;

R^2 is hydrogen or C_1 - C_4 alkyl;

R^3 is hydrogen or C_1 - C_4 alkyl;

25

E, G, L and M are each independently selected from hydrogen and C_1 - C_4 alkyl;

J is hydrogen or C_1 - C_4 alkyl;

30 R^4 is phenyl, mono- or di- or tri-substituted phenyl, pyridyl, thienyl, or furanyl;
wherein the substituents on the substituted phenyl are independently selected from
halogen, cyano, CO_2R^c , OR^c , $\text{CON(R}^c)_2$, methylenedioxy, C_1 - C_4 alkyl and
fluorinated C_1 - C_4 alkyl;

R⁵ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, CO₂R^c, OR^c, CON(R^c)₂, methylenedioxy, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

5

one of R⁶ and R⁷ is hydrogen, and the other of R⁶ and R⁷ is hydrogen or C₁-C₄ alkyl;

- 10 R⁸ is phenyl, mono- or di- or tri-substituted phenyl, pyridyl, or mono- or di- or tri-substituted pyridyl; wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, C₁-C₄ alkyl and CF₃; and wherein the substituents on the substituted pyridyl are independently selected from cyano, nitro, N(R^c)₂, (CH₂)₀₋₄CO₂R^c,
15 (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, phenyl, OR^c, halogen, C₁-C₄ alkyl and CF₃;

R⁹ is hydrogen, C₁-C₄ alkyl, or fluorinated C₁-C₄ alkyl;

- 20 R¹⁰ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c and C₁-C₄ alkyl;

25

R¹³, R¹⁴, R¹⁵ and R¹⁶ are each independently selected from hydrogen and C₁-C₄ alkyl;

R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

30

R^b is C₁-C₄ alkyl or (CH₂)₀₋₃CF₃;

R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^d is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

m, n, o, and p are each independently 0 or 1; and


5 q1 is an integer from 0 to 4;

or a pharmaceutically acceptable salt thereof.

3. The compound according to claim 2, wherein each of A¹, A²,
10 A³ and A⁴ is C-X²;

or a pharmaceutically acceptable salt thereof.

4. The compound according to claim 1, wherein Q is



15

each of A¹, A², A³ and A⁴ is C-X²;

each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl,
20 fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, or (CH₂)₀₋₄OR^a;

each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl,
fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, or (CH₂)₀₋₄OR^a;

25 R¹ is hydrogen, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated
C₃-C₆ cycloalkyl, phenyl, or substituted phenyl, wherein the substituents on the

substituted phenyl are independently selected from halogen, cyano, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl and (CH₂)₀₋₄OR^a;

R² is hydrogen or C₁-C₄ alkyl;

5

R³ is hydrogen or C₁-C₄ alkyl;

R⁴ is phenyl, mono- or di- or tri-substituted phenyl, pyridyl, thienyl, or furanyl; wherein the substituents on the substituted phenyl are independently selected from
10 halogen, cyano, CO₂R^c, OR^c, CON(R^c)₂, methylenedioxy, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

R⁵ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the substituted phenyl are
15 independently selected from halogen, cyano, CO₂R^c, OR^c, CON(R^c)₂, methylenedioxy, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

one of R⁶ and R⁷ is hydrogen, and the other of R⁶ and R⁷ is hydrogen or C₁-C₄ alkyl;

20

R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^b is C₁-C₄ alkyl or (CH₂)₀₋₃CF₃;


25 R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃; and

q₁ is an integer from 0 to 4;

or a pharmaceutically acceptable salt thereof.

30

5. The compound according to claim 4, having the formula:



wherein Z is CH₂, CHOH or C=O;

5 each X³ is independently hydrogen, CF₃, cyano, halogen, or C₁-C₄ alkyl;

R⁵ is hydrogen, cyano, or CO₂R^c;

q₂ is an integer from 0 to 4; and


10

q₃ is an integer from 0 to 3;

or a pharmaceutically acceptable salt thereof.

15

6. The compound according to claim 5, having the formula:



wherein each X^1 is independently hydrogen, fluorine, methyl, ethyl, cyano, CF_3 , methoxy, ethoxy, or OCF_3 ;

5

each X^3 is independently hydrogen, CF_3 , cyano, fluorine, methyl, or ethyl; and

q_1 is an integer from 0 to 2;

10 or a pharmaceutically acceptable salt thereof.

7. The compound according to claim 6, selected from the group consisting of


15 4-(3,4-difluorophenyl)-3-((4-cyano-4-(2-cyanophenyl)piperidin-1-yl)propylcarbamoyl)-3,4-dihydro-quinazolin-2-one;

4-(3,4-difluorophenyl)-3-(4-(4-fluorophenyl)piperidin-1-yl)propylcarbamoyl)-3,4-dihydro-quinazolin-2-one;

20

and pharmaceutically acceptable salts thereof.

8. The compound according to claim 1, wherein Q is



each of A¹, A², A³ and A⁴ is C-X²;

- 5 each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, or (CH₂)₀₋₄OR^a;

each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, or (CH₂)₀₋₄OR^a;

- 10 R¹ is hydrogen, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, fluorinated C₁-C₄ alkyl, fluorinated C₃-C₆ cycloalkyl, phenyl, or substituted phenyl, wherein the substituents on the substituted phenyl are independently selected from halogen, cyano, C₁-C₄ alkyl, fluorinated C₁-C₄ alkyl and (CH₂)₀₋₄OR^a;

- 15 R² is hydrogen or C₁-C₄ alkyl;

R³ is hydrogen or C₁-C₄ alkyl;

- 20 E, G, L and M are each independently selected from hydrogen and C₁-C₄ alkyl;


J is hydrogen or C₁-C₄ alkyl;

- 25 one of R⁶ and R⁷ is hydrogen, and the other of R⁶ and R⁷ is hydrogen or C₁-C₄ alkyl;

R⁸ is selected from phenyl, mono- or di- or tri-substituted phenyl, pyridyl, or mono- or di- or tri-substituted pyridyl; wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, N(R^c)₂, NR^cCOR^c,

- NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, C₁₋₄ alkyl and CF₃; and wherein the substituents on the substituted pyridyl are independently selected from cyano, N(R^c)₂, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c, phenyl, OR^c, halogen, C₁₋₄ alkyl and CF₃;
- 10 R¹⁰ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂, (CH₂)₀₋₄SO₂R^c and C₁₋₄ alkyl;
- 15 R¹³, R¹⁴, R¹⁵ and R¹⁶ are each independently selected from hydrogen and C₁₋₄ alkyl;
- R^a is hydrogen, C₁₋₄ alkyl, or (CH₂)₀₋₃CF₃;
- R^c is hydrogen, C₁₋₄ alkyl, or (CH₂)₀₋₃CF₃;
- 20 R^d is hydrogen, C₁₋₄ alkyl, or (CH₂)₀₋₄CF₃;
- m, n, o, and p are each independently 0 or 1; and
- q₁ is an integer from 0 to 4;
- 25 or a pharmaceutically acceptable salt thereof.

9. The compound according to claim 8, having the formula:



wherein T is C-X³ or N;

- 5 each X³ is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, C₁₋₄ alkyl, or CF₃;

R¹⁰ is hydrogen, cyano, or OR^c;

- 10 R²⁰ is hydrogen or OH;

q₂ is an integer from 0 to 4;

q₃ is an integer from 0 to 3; and

- 15 w is 0 or 1, provided that when w is 0, R²⁰ is hydrogen;

or a pharmaceutically acceptable salt thereof.

- 20 10. The compound according to claim 9, wherein

each X¹ is fluorine;

each X² is hydrogen;

- 25 R²⁰ is hydrogen;

q1 is an integer from 0 to 3; and

w is 1;

5 or a pharmaceutically acceptable salt thereof.

11. The compound according to claim 10, selected from the group consisting of

10 *trans*-4-(3,4-difluorophenyl)-3-(1-(4-(2-cyanophenyl)cyclohexyl)-3-hydroxy-azetidin-3-yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2-one;

trans-4-(3,4-difluorophenyl)-3-(1-(4-(2-cyanophenyl)cyclohexyl)azetidin-3-yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2-one;


15

trans-4-(3,4-difluorophenyl)-3-(1-(4-(2-pyridyl)cyclohexyl)azetidin-3-yl)methylcarbamoyl)-3,4-dihydro-quinazolin-2(3H)-one;

and pharmaceutically acceptable salts thereof.

20

12. The compound according to claim 1, having the formula:



25 wherein

each X^1 is independently hydrogen, halogen, cyano, C_1 - C_4 alkyl, CF_3 , or $(CH_2)_0$ - $4OR^a$;

each X^2 is independently hydrogen, halogen, cyano, C_1 - C_4 alkyl, CF_3 , or $(CH_2)_0$ -
5 $4OR^a$;

each X^3 is independently hydrogen, halogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$,
 $SO_2N(R^c)_2$, SO_2R^c , C_1 - C_4 alkyl, or CF_3 ;

10 R^2 is hydrogen or C_1 - C_4 alkyl;

R^3 is hydrogen or C_1 - C_4 alkyl;

R^9 is hydrogen or C_1 - C_4 alkyl;

15

R^{10} is hydrogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, phenyl, or mono- or di- or tri-
substituted phenyl, wherein the substituents on the phenyl are independently selected
from halogen, cyano, OR^c , $(CH_2)_0$ - $4CO_2R^c$, $(CH_2)_0$ - $4CON(R^c)_2$, $N(R^c)_2$,
 NR^cCOR^c , $NR^cCON(R^c)_2$, $NR^cSO_2R^c$, $NR^cSO_2N(R^c)_2$, $(CH_2)_0$ - $4SO_2N(R^c)_2$,

20 $(CH_2)_0$ - $4SO_2R^c$, C_1 - C_4 alkyl and fluorinated C_1 - C_4 alkyl;

R^a is hydrogen, C_1 - C_4 alkyl, or $(CH_2)_0$ - $3CF_3$;

R^c is hydrogen, C_1 - C_4 alkyl, or $(CH_2)_0$ - $3CF_3$;

25

t is an integer from 2 to 4; and

q_1 , q_2 and q_3 are each independently integers from 0 to 4;

30 or a pharmaceutically acceptable salt thereof.

13. The compound according to claim 12, wherein

each X^1 is independently hydrogen, halogen, cyano, methyl, ethyl, CF_3 , or OCF_3 ;

each X^2 is hydrogen;

each X^3 is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy,
ethoxy, CO_2R^c , $\text{CON}(\text{R}^c)_2$, $\text{SO}_2\text{N}(\text{R}^c)_2$, SO_2R^c , CF_3 , or OCF_3 ;

R^2 is hydrogen;

R^3 is hydrogen;

10

R^9 is hydrogen;

R^{10} is hydrogen, cyano, or OR^c ;

15 R^c is hydrogen, methyl, or ethyl;


t is 2 or 3; and

q_1 and q_3 are each independently integers from 0 to 3;

20

or a pharmaceutically acceptable salt thereof.

14. The compound according to claim 1, having the formula:



25

wherein

each X^1 is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

5 each X^2 is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

each X^3 is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, C₁₋₄ alkyl, or CF₃;

10 R² is hydrogen or C₁-C₄ alkyl;

R³ is hydrogen or C₁-C₄ alkyl;

R⁹ is hydrogen or C₁-C₄ alkyl;

15 R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

20 t is an integer from 2 to 4; and

q₁, q₂ and q₃ are each independently integers from 0 to 4;

or a pharmaceutically acceptable salt thereof.

25

15. The compound according to claim 14, wherein

each X^1 is independently hydrogen, halogen, cyano, methyl, ethyl, CF₃, or OCF₃;

30 each X^2 is hydrogen;

each X^3 is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy, ethoxy, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, CF₃, or OCF₃;

R² is hydrogen;

R³ is hydrogen;

5 R⁹ is hydrogen;

R^c is hydrogen, methyl, or ethyl;

t is 2 or 3; and


10

q₁ and q₃ are each independently integers from 0 to 3;

or a pharmaceutically acceptable salt thereof.

15

16. The compound according to claim 1, having the formula:



wherein

20 each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

25

each X^3 is independently hydrogen, halogen, cyano, OR^c , CO_2R^c , $CON(R^c)_2$, $SO_2N(R^c)_2$, SO_2R^c , C_1-4 alkyl, or CF_3 ;

R^2 is hydrogen or C_1-C_4 alkyl;

5

R^3 is hydrogen or C_1-C_4 alkyl;

R^a is hydrogen, C_1-C_4 alkyl, or $(CH_2)_0-3CF_3$;

10 R^c is hydrogen, C_1-C_4 alkyl, or $(CH_2)_0-3CF_3$;

o and p are each independently 0 or 1;

w is an integer from 0 to 2; and

15

q_1 , q_2 and q_3 are each independently integers from 0 to 4;

or a pharmaceutically acceptable salt thereof.

20

17. The compound according to claim 16, wherein

each X^1 is independently hydrogen, halogen, cyano, methyl, ethyl, CF_3 , or OCF_3 ;

each X^2 is hydrogen;

25

each X^3 is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy, ethoxy, CO_2R^c , $CON(R^c)_2$, $SO_2N(R^c)_2$, SO_2R^c , CF_3 , or OCF_3 ;

R^2 is hydrogen;

30

R^3 is hydrogen;


R^c is hydrogen, methyl, or ethyl;

w is 0 or 1; and

q1 and q3 are each independently integers from 0 to 3;

5 or a pharmaceutically acceptable salt thereof.

18. The compound according to claim 1, having the formula:



10

wherein

each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

15

each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

each X³ is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, C₁₋₄ alkyl, or CF₃;

20

R² is hydrogen or C₁-C₄ alkyl;

R⁹ is hydrogen or C₁-C₄ alkyl;

25

R¹⁰ is hydrogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, phenyl, or mono- or di- or tri-substituted phenyl, wherein the substituents on the phenyl are independently selected from halogen, cyano, OR^c, (CH₂)₀₋₄CO₂R^c, (CH₂)₀₋₄CON(R^c)₂, N(R^c)₂, NR^cCOR^c, NR^cCON(R^c)₂, NR^cSO₂R^c, NR^cSO₂N(R^c)₂, (CH₂)₀₋₄SO₂N(R^c)₂,
5 (CH₂)₀₋₄SO₂R^c, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

10

o and p are each independently 0 or 1;

w is an integer from 0 to 2; and

15

q₁, q₂ and q₃ are each independently integers from 0 to 4;

or a pharmaceutically acceptable salt thereof.

20

19. The compound according to claim 18, wherein

each X¹ is independently hydrogen, halogen, cyano, methyl, ethyl, CF₃, or OCF₃;

each X² is hydrogen;

25

each X³ is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy, ethoxy, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, CF₃, or OCF₃;

R² is hydrogen;

30

R⁹ is hydrogen;

R¹⁰ is hydrogen, cyano, or OR^c;

R^c is hydrogen, methyl, or ethyl;


w is 0 or 1; and

q1 and q3 are each independently integers from 0 to 3;

5

or a pharmaceutically acceptable salt thereof.

20. The compound according to claim 1, having the formula:



wherein

each X¹ is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

15

each X² is independently hydrogen, halogen, cyano, C₁-C₄ alkyl, CF₃, or (CH₂)₀₋₄OR^a;

20 each X³ is independently hydrogen, halogen, cyano, OR^c, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, C₁₋₄ alkyl, or CF₃;

R² is hydrogen or C₁-C₄ alkyl;

25 R⁹ is hydrogen or C₁-C₄ alkyl;

R^a is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

R^c is hydrogen, C₁-C₄ alkyl, or (CH₂)₀₋₃CF₃;

5 o and p are each independently 0 or 1;

w is an integer from 0 to 2; and

q₁, q₂ and q₃ are each independently integers from 0 to 4;

10

or a pharmaceutically acceptable salt thereof.

21. The compound according to claim 20, wherein

15 each X¹ is independently hydrogen, halogen, cyano, methyl, ethyl, CF₃, or OCF₃;

each X² is hydrogen;

20 each X³ is independently hydrogen, halogen, cyano, methyl, ethyl, hydroxy, methoxy, ethoxy, CO₂R^c, CON(R^c)₂, SO₂N(R^c)₂, SO₂R^c, CF₃, or OCF₃;

R² is hydrogen;

R⁹ is hydrogen;

25

R^c is hydrogen, methyl, or ethyl;

w is 0 or 1; and

30 q₁ and q₃ are each independently integers from 0 to 3;

or a pharmaceutically acceptable salt thereof.

22. A pharmaceutical composition comprising the compound according to claim 1 and a pharmaceutically acceptable carrier.

23. A pharmaceutical composition made by combining the
5 compound according to claim 1 and a pharmaceutically acceptable carrier.

24. A process for making a pharmaceutical composition comprising combining the compound according to claim 1 and a pharmaceutically acceptable carrier.
10

25. The composition according to claim 22 further comprising a testosterone 5-alpha reductase inhibitor.

26. The composition according to claim 25, wherein the
15 testosterone 5-alpha reductase inhibitor is a type 1, a type 2, both a type 1 and a type 2, or a dual type 1 and type 2 testosterone 5-alpha reductase inhibitor.

27. The composition according to claim 26, wherein the testosterone 5-alpha reductase inhibitor is a type 2 testosterone 5-alpha reductase
20 inhibitor.

28. The composition according to claim 27, wherein the testosterone 5-alpha reductase inhibitor is finasteride.

25 29. A method of treating benign prostatic hyperplasia in a subject in need thereof which comprises administering to the subject a therapeutically effective amount of the compound according to claim 1.

30 30. The method according to claim 29, wherein the compound does not cause a fall in blood pressure at dosages effective to alleviate benign prostatic hyperlasia.

31. The method according to claim 29, wherein the compound is administered in combination with a testosterone 5-alpha reductase inhibitor.
35

32. The method according to claim 31, wherein the testosterone 5-alpha reductase inhibitor is finasteride.

5 33. A method of treating benign prostatic hyperplasia in a subject in need thereof which comprises administering a therapeutically effective amount of the composition according to claim 22.

34. The method according to claim 33, wherein the composition further comprises a therapeutically effective amount of a testosterone 5-alpha
10 reductase inhibitor.

35. A method of relaxing lower urinary tract tissue in a subject in need thereof which comprises administering to the subject a therapeutically effective amount of the compound according to claim 1.
15

36. The method according to claim 35, wherein the compound is administered in combination with a testosterone 5-alpha reductase inhibitor.

20 37. The method according to claim 36, wherein the testosterone 5-alpha reductase inhibitor is finasteride.

38. A method of treating a condition which is susceptible to treatment by antagonism of the alpha 1a receptor which comprises administering to a subject in need thereof an amount of the compound according to claim 1 effective to
25 treat the condition.

39. A method of eliciting an alpha 1a antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically
30 effective amount of the compound according to claim 1.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/01775**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 544/256, 544/279, 544/286

514/252.16, 514/258, 514/259

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS, MARPAT

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98/57642 A1 (MERCK & CO., INC.) 23 December 1998, see entire document.	1-39
A	WO 98/57640 A1 (MERCK & CO., INC.) 23 December 1998, see entire document.	1-39
A	US 5,389,631 A (CLAREMON et al.) 14 February 1995, see entire document.	1-24

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	
A document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specifying)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O document referring to an oral disclosure, use, exhibition or other means	*A* document member of the same patent family
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 08 MAY 2000	Date of mailing of the international search report 30 MAY 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer Mukund Shah Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/01775

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☒

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/01775

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (7):

C07D 239/82, 487/22, 487/14, 487/04
A61K 31/517, 31/519, 31/496

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

544/256, 544/279, 544/286
514/252.16, 514/258, 514/259

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-39, drawn to the compounds of the formula as depicted in claim 1, their composition and methods of use wherein each of A1, A2, A3, and A4 is C-X2.

Group II, claims 1, 2, 22-39, drawn to the compounds of the formula as depicted in claim 1, their composition and methods of use wherein one of A1, A2, A3, and A4 is N.

Group III, claims 1, 2, 22-39, drawn to the compounds of the formula as depicted in claim 1, their compositions and methods of use wherein A1 and A3 are N, A2 and A4 are C-X2.

Group IV, claims 1, 2, 22-39, drawn to the compounds of the formula as depicted in claim 1, their compositions and methods of use wherein A1 and A4 are N, A2 and A3 are C-X2.

Groups I-IV are directed to structurally dissimilar compounds such that the variable core created by varying the definitions of the formula do not belong to a recognized class of chemical compounds in the art, and references anticipating one invention would not render obvious the others, for example, quinazoline is different from pyridopyrimidine or pteridine, etc. One skilled in the art would not consider such diverse structures as functional equivalents of each other.